



## *Stictis* s.l. (Ostropales, Ascomycota) in the Russian Far East

Eugene S. Popov<sup>1</sup>, Sergey V. Chesnokov<sup>1,2\*</sup>, Liudmila A. Konoreva<sup>2,3</sup>, Alexander K. Ezhkin<sup>4</sup>, Irina S. Stepanchikova<sup>1,5</sup>, Ekaterina S. Kuznetsova<sup>1,5</sup>, Dmitry E. Himelbrant<sup>1,5</sup>, Irina A. Galanina<sup>6</sup> & Svetlana I. Tchabanenko<sup>7†</sup>

Eugene S. Popov<sup>1</sup>  
e-mail: epopov@binran.ru

Sergey V. Chesnokov<sup>1,2\*</sup>  
e-mail: lukinbrat@mail.ru

Liudmila A. Konoreva<sup>2,3</sup>  
e-mail: ajdazapov@yandex.ru

Alexander K. Ezhkin<sup>4</sup>  
e-mail: ezhkin@yandex.ru

Irina S. Stepanchikova<sup>1,5</sup>  
e-mail: stepa\_ir@mail.ru

Ekaterina S. Kuznetsova<sup>1,5</sup>  
e-mail: igel\_kuzn@mail.ru

Dmitry E. Himelbrant<sup>1,5</sup>  
e-mail: d\_brant@mail.ru

Irina A. Galanina<sup>6</sup>  
e-mail: gairka@yandex.ru

Svetlana I. Tchabanenko<sup>7†</sup>

<sup>1</sup> Komarov Botanical Institute RAS, St. Petersburg, 197376, Russia

<sup>2</sup> Botanical Garden-Institute FEB RAS, Vladivostok, Russia

<sup>3</sup> Polar-Alpine Botanical Garden-Institute of the Kola Science Centre RAS, Kirovsk, Russia

<sup>4</sup> Institute of Marine Geology and Geophysics FEB RAS, Yuzhno-Sakhalinsk, Russia

<sup>5</sup> St. Petersburg State University, St. Petersburg, Russia

<sup>6</sup> Federal Scientific Center of East Asian Terrestrial Biodiversity FEB RAS, Vladivostok, 690022, Russia

<sup>7</sup> Sakhalin Branch of Botanical Garden-Institute FEB RAS, Vladivostok, Russia

† Deceased

\* corresponding author

Manuscript received: 08.10.2019

Review completed: 12.02.2020

Accepted for publication: 20.05.2020

Published online: 23.05.2020

### ABSTRACT

This paper summarizes literature data and new information on the geographical distribution and ecology of *Stictis* and closely related genera in Russia. *Stictis sphaeroboloides* is reported as new to Eurasia; *Carestiella socia* and *Stictis populorum* are new to Asia, *S. carnea* is new to Russia, and *S. radiata* is new to the Russian Far East. New data on distribution and ecology of *Carestiella socia* and five *Stictis* species are provided along with a comprehensive key for all species of *Stictis* s.l. sensu Wedin et al. (2006) known in Russia and nearby areas. Our research clearly shows importance of the collaboration between mycologists and lichenologists to study trophically complex taxa and to obtain the most complete knowledge about fungal diversity.

**Keywords:** lichenized fungi, saprotrophic fungi, Sakhalin, Kamchatka, Primorsky Territory, Kuril Islands

### РЕЗЮМЕ

Попов Е.С., Чесноков С.В., Конорева Л.А., Ёжкин А.К., Степанчикова И.С., Кузнецова Е.С., Гимельбрант Д.Е., Галанина И.А., Чабаненко С.И. Род *Stictis* s.l. (Ostropales, Ascomycota) на Российском Дальнем Востоке. Впервые представлен обзор по роду *Stictis* s.l. sensu Wedin et al. (2006) для российского Дальнего Востока. Составлен ключ для видов, известных в России. Представлены новые данные о распространении и экологии *Carestiella socia* и пяти видов из рода *Stictis*. *Stictis sphaeroboloides* впервые приводится для Евразии. *Carestiella socia* и *Stictis populorum* являются новыми видами для Азии; *S. carnea* впервые приводится для России, *S. radiata* приводится здесь как новый вид для российского Дальнего Востока. Наше исследование ясно показывает важность сотрудничества между микологами и лишенологами для изучения трофически сложных таксонов и получения наиболее полных знаний о разнообразии грибов.

**Ключевые слова:** лишенизированные грибы, сапротрофные грибы, Сахалин, Камчатка, Приморский край, Курильские острова

*Stictis* s.l. as circumscribed by Wedin et al. (2006) includes taxa from at least three ostropalean genera (*Stictis* s.str., *Carestiella*, and *Schizoxylon* pro parte). Within *Stictis* s.l. some taxa have shown facultative lichenization where the fungus may either persist as saprotroph or be lichenized, apparently depending on its substrate, i. e. bark or wood (Wedin et al. 2004).

Four *Schizoxylon* species, five *Stictis* species and one *Ostropa* species were reported from Russia before. Their distribution by large geographical regions (adopted after Czerepanov 1995) is as follows:

**European Part of Russia.** Five species of *Stictis* and one species of *Schizoxylon* are known: *Schizoxylon berkeleyanum* (Durieu et Lév.) Fuckel (Jaczewski 1896), *Stictis brunnescens* Gilenstam et al. (Wedin et al. 2006, Notov et al. 2011, Kuznetsova et al. 2012), *S. mollis* Pers. (Jaczewski 1895, Tranzschel 1905, Tranzschel & Serebrianiokov 1912, Popov 2012, Muchnik et al. 2019), *S. phragmitis* Lobik (Lobik 1928), *S. populorum* (Gilenstam) Gilenstam (Hermansson et al. 2006, Urbanavichus 2014), *S. radiata* Pers. (Martius 1817, Weinmann 1828, Karsten 1866, Vasil'eva 1939, Sirko 1976, Vaasma et al. 1986, Urbanavichene 2011, Kuznetsova et al. 2012, Popov & Volobuev 2014, Churakov et al. 2015, Huseyin et al. 2016, Himelbrant et al. 2017).

**Caucasus.** Three species were previously reported: *Schizoxylon alboatrum* (Urbanavichus & Urbanavichene 2017), *Stictis brunnescens* Gilenstam et al. (Urbanavichus & Urbanavichene 2015), and *S. radiata* Pers. (Vaasma et al. 1986, Urbanavichene & Urbanavichus 2014).

**Western and Eastern Siberia.** Only *Stictis mollis* has been known to occur in Siberia since 1880 (Thümen, as *S. ollaris* Wallr.). Konoreva et al. (2016) summarized data on the distribution and ecology of *Stictis* s.l. in Russia and added *Schizoxylon albescens* Gilenstam et al., *Stictis brunnescens* and *S. radiata* from Altai Mts. and Eastern Siberia.

**Russian Far East.** Three species were previously reported: *Stictis mollis*, *Schizoxylon berkeleyanum* and *Ostropa barbara* (Koval' 1972) from the Primorye Territory, and *Schizoxylon hemisphaericum* (Fr.) DiCosmo et al. known only from the type locality in Kamchatka (Fries 1823, DiCosmo et al. 1984).

Thus, most of known records originate from European part of Russia, while the diversity and distribution of Stictidaceae in Asian part remain poorly understood, and only four species were known from the Russian Far East before our research.

## MATERIAL AND METHODS

The present study is based on the specimens collected by the authors between 2005 and 2018 in various regions of the Russian Far East. The data from the Pskov Region and Karachayevo-Circassian Republic were used to clarify the distribution of the species in Russia. The specimens are deposited in the lichen and fungal subdivisions of the herbarium of the Komarov Botanical Institute RAS, St. Petersburg, Russia (LE), also in the herbaria of St. Petersburg State University, St. Petersburg, Russia (LECB), Polar-Alpine Botanical Garden-Institute of the KSC RAS, Kirovsk, Russia (KPABG), University of Helsinki, Helsinki, Finland (H), Institute of Marine Geology and Geophysics FEB RAS, Yuzhno-Sakhalinsk, Russia (SAK) and Sakhalin Branch of Botanical Garden-Institute FEB RAS, Yuzhno-Sakhalinsk, Russia (SAKH).

The material was examined at The Core Facility Center "Cell and Molecular Technologies in Plant Science" at the Komarov Botanical Institute using standard microscopic techniques (Smith et al. 2009). Photographs of the species were made using AxioCam MRC5 digital camera mounted on Stemi-2000 CS light microscope. The distribution map was prepared using MapInfo GIS software. Geographical coordinates are given in spatial reference system WGS 1984.

Abbreviations of authors in text: EP = Eugene S. Popov, SCh = Sergey V. Chesnokov, LK = Liudmila A. Konoreva, AE = Alexander K. Ezhkin, IS = Irina S. Stepanchikova, EK = Ekaterina S. Kuznetsova, DH = Dmitry E. Himelbrant, IG = Irina A. Galanina.

## RESULTS

Altogether 21 specimens were studied, and 7 species of *Stictis* s.l. were revealed in the Russian Far East. *Carestiella socia*, *Stictis popolorum*, and *S. sphaeroboloides* are new to Asia, *S. carnea* is new to Russia, and *S. radiata* is new to the Russian Far East.

To facilitate the progress in the studies of diversity and distribution of Stictidaceae in Russia we provide an identification key for the filiform-spored epixylic and epiphytic species of *Stictis* and morphologically similar genera known to occur in the country. Short descriptions, differences from the related species, notes on ecology and distribution are given for species found in the Far East.

## Key to the epixylic and epiphytic filiform-spored species of Stictidaceae in Russia

All species mentioned in the key are known from Russia; for the species which are not yet reported from Russian Far East but expected to be found there the notes on distribution are given in brackets.

1. Apothecia opening at maturity by a rounded pore or irregular fissures ..... 2
  - Apothecia opening at maturity by a transverse slit ..... 13
2. Periphysoids  $\pm$  distinct; apothecial margin with distinct crystalline inclusions; ascospores not disarticulating into part-spores (genus *Stictis*) ..... 3
  - Periphysoids absent or indistinct; apothecial margin usually without crystalline inclusions (but see *Schizoxylon alboatrum*), sometimes with crystalline pruina on outer surface; ascospores disarticulating or not ..... 9
3. Hymenium remaining covered by crystalline layer for a long time after eruption through the host tissues ..... *Stictis sphaeroboloides*
  - Hymenium soon opening by a pore ..... 4
4. Disc ochraceous, glossy, never pruinose ..... 5
  - Disc brown, or grey to black,  $\pm$  dull, often pruinose ..... 7
5. Margin partly brown in section ..... *Stictis brunnescens*
  - Margin colourless in section ..... 6
6. Ascospores 3–3.5  $\mu$ m broad, distinctly tapered below; individual cells 3–4  $\mu$ m long ..... *Stictis carnea*
  - Ascospores 2–2.5  $\mu$ m broad, not or slightly tapered; individual cells 4–7  $\mu$ m long ..... *Stictis radiata*
7. Ascospores more than 3  $\mu$ m broad ..... *Stictis popolorum*
  - Ascospores less than 3  $\mu$ m broad ..... 8
8. Ascospores longer than 100  $\mu$ m; on wood and bark ..... *Stictis mollis*
  - Ascospores shorter than 100  $\mu$ m; on stems of *Phragmites* ..... *Stictis phragmitis*
    - [so far known only from the type locality in Kalmykia]
9. Ascospores with average length less than 100  $\mu$ m, 3- to 7-septate; asci less than 150  $\mu$ m long, multisporous ..... *Carestiella socia*
  - Ascospores and asci longer on average; ascospores with more than 10 septa; asci initially 8-spored, when mature filled with numerous part-spores ..... 10
10. Ascospores 100–130  $\times$  5–6  $\mu$ m, part-spores 1-celled, globose to discoid, 3–7  $\times$  5–6  $\mu$ m; apothecia black ..... *Schizoxylon hemisphaericum*
  - Ascospores more than 150  $\mu$ m long and up to 2.5  $\mu$ m broad, part-spores, if present, 1- to multiseptate, oblong; apothecia with grayish, white or yellowish pruina ..... 11
11. Excipular pigment dissolving in KOH and staining the medium bright yellow; ascospores regularly disarticulating into two-celled part-spores; predominantly on large herbaceous stems ..... *Schizoxylon berkeleyanum*
  - Excipular pigment not reacting with KOH; ascospores disarticulating or not; on wood and bark ..... 12
12. Ascospores ca 150–270  $\times$  1.5–2.5  $\mu$ m, disarticulating into 2–15-septate part-spores; asci ca 200–280  $\mu$ m; margin without crystalline inclusions ..... *Schizoxylon albescens*
  - [Palearctic taxon]
  - Ascospores longer, not regularly disarticulating into part-spores; asci longer than 300  $\mu$ m; margin with crystals ..... *Schizoxylon alboatrum* [Holarctic taxon]
13. Apothecia deep immersed, hockey stick-shaped in longitudinal section, with longer axis subparallel to substrate surface ..... *Robergea cubicularis* [Holarctic taxon]
  - Apothecia erumpent,  $\pm$  rounded in longitudinal section, oriented perpendicular to substrate surface ..... *Ostropa barbara*




Figure 1 A – *Carestiella socia* Bres.; B – *Stictis carnea* Seaver & Waterston. Scale bar A, B = 1 mm

***Carestiella socia*** Bres., Malpighia 11(6–8): 274. 1897 (Fig. 1A, 2)

This species is easily recognized by its multisporous asci, relatively short, often falcate or sigmoid ascospores with 3–7 transverse septa, and peculiar structure of the excipular margin, which is composed of loosely interwoven brown thick-walled hyphae without crystalline inclusions. The only known specimen of *C. socia* in Russia was collected during Ryabushinsky's Kamchatka Expedition (1908–1909), but remained unidentified and was not mentioned in the expedition report (Tranzschel 1914). It fits well the detailed description of the species given by Wedin et al. (2006). In the material studied asci were predominately 16-spored, ca 80–90 × 10–15 μm. Ascospores 38–50 × 2–2.5 μm, with 3–7 septa.

**Ecology.** Saprotrophic on wood and bark of *Populus* and *Salix* spp., facultatively lichenized according to Wedin et al. (2006).

**Distribution.** This is the first report of *Carestiella socia* from Russia and Asia. Previously the species was reported from Europe (Italy, Sweden, Norway), and North America (Wedin et al. 2006).

**Specimens examined:** KAMCHATKA TERRITORY, Yelizovo District, between Malka and Ganaly, near the Shestaya River, on bark of *Salix* sp. [according to the label on 'vetla', a Russian name for some species of *Salix* subgen. *Salix*], VI.1909, V. P. Savicz s. n. (LE 172078).

***Schizoxylon hemisphaericum*** (Fr.) DiCosmo, Nag Raj & W.B. Kendr., Can. J. Bot. 61(1): 44. 1983

*Schizoxylon hemisphaericum* was described by Fries (1823) as *Phacidium hemisphaericum* Fr. based on single collection made by Wormskjold in Kamchatka. DiCosmo et al. (1984) studied the holotype which is kept now in Fries's Herbarium in Uppsala (UPS) and argued that this species should be placed in *Schizoxylon*. They also provided a full description and illustrations of the type material.

**Ecology.** Saprotrophic on bark and wood of *Betula*.

**Distribution:** Known only from the type locality.

***Stictis carnea*** Seaver & Waterston, Mycologia 33(3): 311. 1941 (Fig. 1B, 2)

*Stictis carnea* has apothecia with a general appearance similar to those of *S. radiata*, but differs in definitely broader ascospores




Figure 2 The distribution of genera *Stictis* and *Carestiella* in the Russian Far East: 1 – *Stictis radiata* Pers., 2 – *S. mollis* Pers., 3 – *S. populorum* (Gilenstam) Gilenstam, 4 – *S. carnea* Seaver & Waterston, 5 – *S. sphaeroboloides* (Schwein.) Ellis, 6 – *Carestiella socia* Bres.

with short, isodiametric to discoid individual cells. For a full description and illustrations, refer to Sherwood (1977).

**Ecology.** Saprotrophic on various woody substrates.

**Distribution.** Widely distributed in tropical areas. Known from Asia (Hong Kong, Philippines, Sri Lanka), Africa, South America, New Zealand (Sherwood 1977, Johnston 1983, Whitton et al. 1999). The record from this survey is the most northern known occurrence of the species (Fig. 2). This, however, is in good agreement with the relatively high diversity of taxa with tropical and subtropical biogeographic affinities in the flora of the southern Primorye Region of the Russian Far East (Kozhevnikov & Kozhevnikova 2014).

**Specimens examined:** PRIMORYE TERRITORY, Khasansky District, Kedrovaya Pad Nature Reserve, the Kedrovaya River valley, 43°06'00.5"N 131°33'32.3"E, on bark of *Acer* sp. 19.VIII.2005, EP (LE 323496).

***Stictis mollis*** Pers., Mycol. eur. (Erlanga) 1: 337. 1822 (Fig. 2)

*Stictis mollis* is one of the three *Stictis* species (together with *S. confusum* and *S. populorum*), shown to occur on wood as a saprotroph, and on bark when lichenized (Wedin et al. 2006).

*Stictis mollis* is easily distinguished from the two other *Stictis* species with distinctly brown margins (*S. confusum* and *S. populorum*) by the comparatively narrow spores and the olivaceous, amorphous pigmentation of the uppermost part of the paraphyses (Wedin et al. 2006).

**Ecology.** According to Wedin et al. (2006) the saprotrophic form of the species inhabits decorticated thin dead branches or twigs of *Populus tremula* L. and *Salix caprea* L. in Sweden, Norway and Germany, while the lichenized form is usually developed on young, thin smooth-barked *Populus*-trunks in Sweden and Norway. The specimen reported by Muchnik et al. (2019) belongs to the lichenized form and grows on bark of young *Acer* sp. The two specimens examined as part of this study were collected on old, coarse bark of *Populus maximowiczii* Henry in mixed forests, and were not lichenized.

**Distribution.** The species is known from Greenland (Rostrup 1891), Norway, Sweden, Germany (Wedin et al. 2006), Spain (Malençon & Bertault 1972), Greece (Zervakis et al. 1999), Ukraine (Dudka et al. 2009), and North Africa (Malençon & Bertault 1969). In Russia it is known from the Ryazan Region (Muchnik et al. 2019), Novgorod Region (Popov 2012), Smolensk Region (Jaczewski 1895), Yaroslavl Region (Tranzschel & Serebriankow 1912), Republic of Crimea (Tranzschel 1905), Krasnoyarsk Territory (Thümen 1880), Primorye Territory (Koval' 1972), Pskov Region, Karachayevo-Circassian Republic (this paper).

**Specimens examined:** SAKHALIN REGION, Sakhalin Island, Korsakov District, Tunaicha Lake surroundings, 46°48'07.4"N 143°06'15.7"E, mixed forest, on bark of *Populus maximowiczii*, 26.X.2017, AE (SAK 1675, LE); *ibid.*, Nevelskiy District, Shebunino village surroundings, 46°26'09.2"N 141°54'12.5"E, alt. 13 m, mixed forest, on bark of *Populus maximowiczii*, 17.V.2016, AE (SAK 1676, LE).

**Additional specimens examined:** PSKOV REGION, Loknyanskiy District, Bashovo, 56°39'31.0"N 30°10'15.7"E, on bark of *Ulmus laevis* Pall., 17.VIII.2011,

EP (LE 247491); *ibid.*, Skokovo, 56°39'45.9"N 30°08'50.5"E, on wood of *Populus longifolia* Fisch. ex Loudon, 28.VII.2011, EP (LE 247555); Ivantsevo, 56°38'47.4"N 29°54'38.7"E, on wood of *Salix caprea* L., 13.VIII.2011, EP (LE 247493); *ibid.*, Polistovsky Nature Reserve, Korolyova Borina 9 km N of Gogolevo, 57°04'40.5"N 30°32'52.0"E, on wood of *Populus tremula* L., 19.IX.2018, EP (LE 323495). KARACHAYEVO-CIRCASSIAN REPUBLIC, Karachayevsky District, Teberdinsky Nature Reserve, environs of Dombay, Amnauz valley, 43°18'26.6"N 41°38'04.1"E, on bark of *Salix caprea* L., 14.VIII.2012, EP (LE 247819).

***Stictis populorum*** (Gilenstam) Gilenstam, Lichenologist 37(1): 74. 2005 (Fig. 2)

A detailed description of this species and differences from the closely related species was given by Wedin et al. (2006).

**Ecology.** A saprotroph on *Populus* wood, particularly on decorticated branches still remaining on the living trunks, or as a lichen in and around cracks in the bark of *Populus* spp. in northern Scandinavia (Wedin et al. 2006). Kamchatian specimen were found on twigs of *P. tremula* in the young spruce forest.


**Distribution.** In Russia this species was reported from the Murmansk Region (Urbanavichus 2014) and Republic of Komi (Hermansson et al. 2006). Originally known from Sweden (Wedin et al. 2006).

**Specimen examined:** KAMCHATKA TERRITORY, Eastern Kamchatka, Mil'kovo District, Kronotsky Nature Reserve, Levaya Shchapina River basin, right bank of the Levaya Shchapina River opposite to the mouth of the Ipuin River, on the hill, 55°07'04"N 159°57'41"E, 280 m, young spruce (*Picea jezoensis* Carrière subsp. *ajanensis* (Fisch. ex Carrière) Silba) forest with sparse grass and traces of burn, on dead twig of *Populus tremula*, 14.VIII.2009, DH, IS, K-15-09 (H).

***Stictis radiata*** Pers., Observ. mycol. (Lipsiae) 2: 73. 1800 (Fig. 2)

A detailed description of the species and differences from the closely related taxa was provided by Wedin et al. (2006).

**Ecology.** The species has wide ecological niche (Wedin et al. 2006). In the Far East it is distributed in different types of forests, mainly near the water (along rivers, creeks and



**Figure 3** The apothecia of *Stictis sphaeroboloides* (Schwein.) Ellis. Scale bar = 1 mm.

lakes), on bark or wood of *Picea*, *Pinus*, *Populus*, *Ulmus*, *Salix* and *Sorbus* spp.

**Distribution.** A rather common species in Russia, *S. radiata* was reported from Murmansk Region (Karsten 1866, Konoreva et al. 2016), Leningrad Region (Kuznetsova et al. 2012, Himelbrant et al. 2017), St. Petersburg (Weinmann 1828), Moscow (Martius 1817), Ulyanovsk Region (Churakov et al. 2015, Huseyin et al. 2016), Kaluga Region (Popov & Volobuev 2014), Chelyabinsk Region (Urbanavichene 2011), Sverdlovsk Region (Sirko 1976), Krasnodar Territory (Vasil'eva 1939, Urbanavichene & Urbanavichus 2014), Republic of Adygea (Vaasma et al. 1986), Krasnoyarsk Territory, Trans-Baikal Territory and Yakutia (Konoreva et al. 2016), Karachayevo-Circassian Republic (this paper). Cosmopolitan species widely distributed in Eurasia, Africa, North and South America Australia, and New Zealand (Rodway 1924, Sherwood 1977, Johnston 1983, Gamundí et al. 2004).

**Specimens examined:** KAMCHATKA TERRITORY, Eastern Kamchatka, Mil'kovo District, Kronotsky Nature Reserve, Levaya Shchapina River basin, 2.3 km SE to the mouth of the Ipuin River, right bank of the Ipuin River, 55°06'05"N 159°59'22"E, 280 m, *Populus suaveolens* Fisch. and graminoid dominated flood-plain forest near the river, on branch of *Picea jezoensis* subsp. *ajanensis*, 10.VIII.2009, DH, IS, K-10-09 (H); *ibid.*, SW part of Askhachny Ridge, 2.2 km N to the mouth of the Ipuin River, SSE slope, 55°08'07"N 159°57'23"E, 300 m, spruce (*Picea jezoensis* subsp. *ajanensis*) forest with green mosses, on branch of *Picea jezoensis* subsp. *ajanensis*, 11.VIII.2009, DH, IS, K-11-09 (H). PRIMORYE TERRITORY, Terney District, Sikhote-Alin, valley of the Taratay creek, 45°42'00.4"N 136°39'48.6"E, *Pinus koraiensis* Siebold et Zucc. dominated forest mixed with broadleaved trees, on wood, 13.08.2010, EK, Prim-22 (LECB); *ibid.*, Krasnoarmeysky District, neighborhood of the Tayozhny village, 45°41'22.6"N 136°13'12.5"E, cedar-spruce forest on the northern slope, on bark of *Picea* sp., 06.VII.2011, IG, T. 412 (KRABG); KHABAROVSK TERRITORY, Khabarovsk District, Bolshekhkhtsirsky Reserve, Bykov River, neighborhood of the cordon Bychikha, 48°14'17.3"N 134°48'49.8"E, 283 m, coniferous-deciduous forest, on branch of *Picea* sp., 29.IX.2018, Sch (LE); *ibid.*, neighborhood of the cordon Sosninsky, Sosninsky creek, 48°14'16.8"N 134°46'41.7"E, 408 m, *Ulmus* sp.–*Betula* sp. forest with *Picea* sp., on branch of *Picea* sp., 01.X.2018, Sch (LE). SAKHALIN REGION, Sakhalin Island, Dolinsky District, neighborhood of the Sokol settlement, Belaya River, 47°15'14.5"N 142°48'21.9"E, 99 m, *Salix* sp. flood-plain forest, on bark of *Salix* sp., 08.V.2017, LK (LE); *ibid.*, 47°15'13.0"N 142°48'12.0"E, 87 m, on wood of *Salix* sp., 08.V.2017, Sch (LE); *ibid.*, 47°15'00.3"N 142°47'33.3" E, alt. 56 m, riparian forest, bark of *Salix udensis* Trautv., 08.V.2016, AE (SAK 1301, 1303, 1304, 1305); *ibid.*, Yuzhno-Sakhalinsk city surroundings, Mitsul' Mountain, 47°03'02.3"N 142°30'39.5"E, alt. 536 m, dark coniferous forest with *Betula ermanii* Cham., on bark of *Sorbus commixta* Hedl., 26.V.2012, AE (SAK 1296); *ibid.*, Tomarinsk District, near Ainskoe Lake, 48°27'03.7"N 142°04'01.6"E, alt. 4 m, mixed forest, on bark of *Picea jezoensis* (Siebold & Zucc.) Carrière, 15.IX.1997, ST (SAKH); *ibid.*, Tymovskiy District, Pilenga River,

51°04'52.4"N 142°44'14.5"E, alt. 119 m, riparian forest, bark of *Populus maximowiczii*, 04.VI.2017, AE (SAK 1297, 1298, 1299); *ibid.*, 51°01'45.7"N 142°50'33.7"E, alt. 154 m, riparian forest, on bark of *Ulmus laciniata* Trautv., 05.VI.2017, AE (SAK 1302); Iturup Island, Ostrovnoy Reserve, Odessky Bay, 44°47'58.7"N 147°13'46.5"E, 17 m, *Abies* sp. forest with *Sasa kurilensis* (Rupr.) Makino & Shibata, on decayed wood of *Sorbus* sp., 13.VIII.2017, LK (LE); Kunashir Island, proximity of Lagunoe Lake, 44°02'50.2"N 145°46'01.6"E, alt. 79 m, old-growth mixed coniferous/broadleaved forest, on bark of *Ulmus laciniata*, 08.I.2017, AE (SAK 1300).

**Additional specimens examined:** KARACHAYEVO-CIRCASSIAN REPUBLIC, Karachayevo District, Teberdinsky Nature Reserve, environs of Dombay, Amnauz Valley, 43°18'26.6"N 41°38'04.1"E, on coniferous wood, 14.VIII.2012, EP (LE 247817); *ibid.*, on wood of *Picea orientalis* (L.) Peterm., 14.VIII.2012, EP (LE 247818).

*Stictis sphaeroboloidea* (Schwein.) Ellis, N. Amer. Fung., Ser. 1: no. 463. 1881 (Fig. 2, 3)

This species is well characterized by erumpent apothecia that remain closed by crystalline layer of ascomatal wall for a long time, and paraphyses with inflated brown apices.

**Ecology.** Saprotrophic on wood and bark of deciduous trees and shrubs, typically on *Quercus* (Sherwood 1977).

**Distribution.** The species was previously known from eastern North America, where it is quite common (Sherwood 1977). The record in Russian Far East thus is another example of a 'Grayan disjunction' between eastern Asia and eastern North America, which is well known in plants (Li 1952, Graham 1972, Davidse 1983, Wen 1999), and not uncommon in fungi (Culberson 1972, Wu & Mueller 1997, Kovacs et al. 2008, Vasilyeva & Stephenson 2010).

**Specimens examined:** PRIMORYE TERRITORY, Khasansky District, Kedrovaya Pad' Nature Reserve, the Kedrovaya River valley, near the mouth of the 1st Zolotoy creek, 43°06'31.9"N 131°31'59.3"E, on bark of *Fraxinus mandshurica* Rupr., 20.VIII.2005, EP (LE 323498).

## DISCUSSION

Symbiotic associations with photosynthetic partners most probably were established and lost independently multiple times in different clades of Ascomycota (Hawksworth 2015), thus many ascomycete taxa include both lichenized and nonlichenized species. The family Stictidaceae is one of the textbook examples of this complexity. It accommodates 54 lichenized species from 13 genera (Lücking et al. 2016) and a number of nonlichenized (Wedin et al. 2005). The level of lichenization in the family varies widely not only at the genus level, but also at the species level, i. e. lichenized and non-lichenized individuals can be found even within one fungal species. This results in lack of attention or neglecting the whole family or certain genera either by lichenologists or by mycologists. In traditional scientific practice the collections of lichenized and nonlichenized species from the same families or even genera are kept in different herbarium compartments (lichenological and mycological, respectively), many specialists poorly know 'alien taxa', 'alien herbaria', and 'alien literature' or even are not interested in 'mixed taxa'. Predictable result of this 'independence' is the

producing of ‘new records’, already successfully published before, as well as lack of our knowledge on such groups.

Our present paper summarizes data on known diversity of the closely related (Wedin et al. 2005) genera *Carestiella*, *Schizoxylon*, and *Stictis* in Russian Far East. Some data are based on recent collections of the authors (*Stictis carnea*, *S. mollis*, *S. populorum*, *S. radiata*, *S. sphaeroboloides*), others are received from the revision of the herbarium samples (*Carestiella socia*), or previously published in mycological papers, like *Schizoxylon hemisphaericum* (DiCosmo et al. 1984). However, Russian Far East is a huge territory, which still remains underexplored, so far we expect that other species can be found there in course of further research. Therefore, our key includes all the species known in Russia at present time, and hopefully can be used in wider territory than Far East only.

The species discussed here are known to have different level of lichenization: saprotrophic – *Schizoxylon hemisphaericum* (DiCosmo et al. 1984), *Stictis carnea* (Sherwood 1977), *S. radiata* (Wedin et al. 2006), and *S. sphaeroboloides* (Sherwood 1977); saprotrophic, but possibly facultatively lichenized depending on a substrate – *Carestiella socia* (Wedin et al. 2006); saprotrophic and clearly lichenized depending on a substrate – *Stictis mollis* (Wedin et al. 2006) and *S. populorum* (Wedin et al. 2006). However, there are still a lot of questions to the trophical strategies of these taxa to be solved using special methods. For instance, it is not clear yet, if the lichenized Stictidaceae, use photobiont as the only source of nutrients, or they get organic compounds also from the substrate (e. g. bark of tree).

Our research clearly shows importance of the collaboration between mycologists and lichenologists to study trophically complex taxa and to gain a better understanding of fungal diversity.

## ACKNOWLEDGEMENTS

The work was carried out in the frame of the institutional research projects no. AAAA-A19-119020690077-4 (work by SCh, LK, IS, EK, DH) and no. AAAA-A19-119020890079-6 (EP) of the Komarov Botanical Institute RAS, no. AAAA-A17-117062710098-4 (IG) of the Federal Scientific Center of East Asian Terrestrial Biodiversity FEB RAS, and the project ‘Cryptogamic biota of Pacific Asia: taxonomy, biodiversity, species distribution’ of the Botanical Garden-Institute FEB RAS (SCh, LK, AE). The study was partly supported by the Russian Foundation for Basic Research (grant no. 18-04-00098, AE; grant no. 18-05-60093 SCh, DH, LK, EK, IS) and no. 19-54-50010 RFBR together with Japan Society for the Advancement of Science; SCh, AE, IG).

## LITERATURE CITED

Churakov, B.P., E.S. Huseyin, F. Selcuk, K.E. Kornilin & T.A. Romanova 2015. Preliminary synopsis of biota of micromycetes of forests of the Ulyanovsk region. In: *Biodiversity and ecology of fungi and fungus-like organisms northern Eurasia* (A.V. Mukhin, ed.), pp. 273–276, Yekaterinburg, publishing house of the Ural University (in Russian). [Чураков Б.П., Хусейн Э.С., Сельчук Ф., Корнилин К.Е., Романова Т.А. 2015. Предварительный конспект биоты микромицетов лесов Ульяновской области // Биоразнообразие и экология грибов и грибоподобных организмов северной Евразии. Екатеринбург: Издательство Уральского университета. С. 273–276].

- Culberson, W.L. 1972. Disjunctive distributions in the lichen-forming fungi. *Annals of the Missouri Botanical Garden* 59(2): 165–173.
- Czerepanov, S.K. 1995. *Vascular plants of Russia and adjacent states (The Former USSR)*. Cambridge, New York, Melbourne, Cambridge University Press, 516 pp.
- DiCosmo, F., T.R. Nag Rag & W.B. Kendrick 1984. A revision of the Phacidaceae and related anamorphs. *Mycotaxon* 21:1–234.
- Davidse, G. 1983. Biogeographical relationships between temperate eastern Asia and temperate eastern North America: the twenty-ninth annual systematics symposium. *Annals of the Missouri Botanical Garden* 70:421–422.
- Dudka, I.O., V.P. Heluta, T.V. Andrianova, V.P. Hayova, Yu. Ya. Tykhonenko, M.P. Prydiuk, Yu.I. Holubtsova et al. 2009. *Fungi of the nature reserves and national nature parks of Eastern Ukraine, vol. 1*. Aristey, Kyiv, 306 pp. (in Ukrainian). [Дудка І.О., Гелюта В.П., Андрианова Т.В., Гайова В.П., Тихоненко Ю.Я., Придюк М.П., Голубцова Ю.І. и др. 2009. Гриби заповідників та національних природних парків Львівської України. Київ: Арістей. Т. 1. 306 с.]
- Fries, E.M. 1823. *Systema Mycologicum, sistens fungorum ordines, genera et species, huc usque cognitae qas ad normam methodi naturalis determinavit, disposuit atque descripsit*. *Lundae: ex officina Berlingiana* 2(2):275–621.
- Gamundí, I.J., D.W. Minter, A.I. Romero, V.A. Barrera, A.L. Gaiotti, M.I. Messuti & M. Stecconi 2004. Checklist of the Discomycetes (Fungi) of Patagonia, Tierra del Fuego and adjacent Antarctic areas. *Darwiniana* 42(1–4):63–164.
- Graham, A. 1972. Outline of the origin and historical recognition of floristic affinities between Asia and eastern North America. In: *Floristics and Paleofloristics of Asia and Eastern North America* (Graham, A., ed.), pp. 1–18. Elsevier, Amsterdam.
- Hawksworth, D. 2015. Lichenization: The origins of a fungal life-style. In: *Recent advances in lichenology: Modern methods and approaches in lichen systematics and culture techniques, vol. 2*, (Upreti, D.K., Divakar, P.K., Shukla, V., Bajpai, R., eds), pp. 1–10. Springer, New Delhi.
- Hermansson, J., T.N. Pystina, B. Ove-Larsson & M.P. Zhurbenko 2006. Lichens and lichenicolous fungi of the Pechoro-Ilychsky Nature Reserve. *Flora i fauna zapovednikov* 109: 1–79 (in Russian). [Херманссон Я., Пыстина Т.Н., Ове-Ларссон Б., Журбенко М.П. 2006. Лишайники и лишенофильные грибы Печеро-Ильчского заповедника // Флора и фауна заповедников. Вып. 109. С. 1–79].
- Himelbrant, D.E., I.S. Stepanchikova, J. Motiejūnaitė, J.V. Gerasimova, E.S. Kuznetsova, A.V. Dyomina & A.G. Tsurykau 2017. New records of lichens and allied fungi from the Leningrad Region, Russia. VIII. *Folia Cryptogamica Estonica* 54:63–70.
- Huseyin, E.S., F. Selcuk, B.P. Churakov, K.E. Kornilin & T.A. Romanova 2016. Microfungi on forest trees and shrubs of Duzce Province (Turkey) and Ulyanovsk Region (Russia). *Mikologiya i Fitopatologiya* 50(1):35–42 (in Russian with English summary). [Хусейн Э.С., Сельчук Ф., Чураков Б.П., Корнилин К.Е., Романова Т.А. 2016. Микромицеты деревьев и кустарников лесов провинции Дюздже (Турция) и Ульяновской области (Россия) // Микология и фитопатология. Т. 50, вып. 1. С. 35–42].
- Jaczewski, A.A. 1895. A catalog of mushrooms from Smolensk province, collected in 1892 and 1894. *Bulletin de la Société Impériale des Naturalistes de Moscou* 9:128–148 (in Russian). [Ячевский А.А. 1895. Каталог грибов Смоленской губернии, собранных в 1892 и 1894 годах // Bulletin de la Société Impériale des Naturalistes de Moscou. Т. 9. С. 128–148].

- Jaczewski, A. 1896. III Série de Matériaux pour la Flore Mycologique du gouvernement de Smolensk. *Bulletin de la Société Impériale des Naturalistes de Moscou* 10(1):65–94.
- Johnston, P.R. 1983. Stictis and its anamorphs in New Zealand. *New Zealand Journal of Botany* 21:249–279.
- Karsten, P.A. 1866. Enumeratio fungorum et myxomycetum in Lapponia orientali aestate 1861 lectorum. *Notiser ur Sällskapetets pro Fauna et Flora Fennica förhandlingar* 8:193–224.
- Konoreva, L.A., S.V. Chesnokov & E.A. Davydov 2016. Stictis and *Schizoxylon* (Stictidaceae, Ostropales) in Russia. *Herzogia* 29(2):706–711.
- Kovacs, G.M., J.M. Trappe, A.M. Alsheikh, K. Bóka & T.F. Elliott 2008. Imaia, a new truffle genus to accommodate *Terfezia gigantea*. *Mycologia* 100(6):930–939.
- Koval', E.Z. 1972. Mycoflora of the reserve of "Kedrovaja pad". In: *Flora and vegetation of the reserve "Kedrovaja pad"* (P.G. Gorovoi, ed.), pp. 105–144 (in Russian). [Коваль Э.З. 1972. Микофлора заповедника «Кедровая Падь» // Флора и растительность заповедника «Кедровая Падь» / под ред. П.Г. Горового. С. 105–144].
- Kozhevnikov, A.E. & Z.V. Kozhevnikova 2014. Taxonomic composition and special features of the natural flora in the Primorsky Territory. *V.L. Komarov Memorial Lectures* 62:7–62 (in Russian with English summary). [Кожевников А.Е., Кожевникова З.В. 2014. Таксономический состав и особенности природной флоры Приморского края // Комаровские чтения. Вып. 62. С. 7–62].
- Kuznetsova, E.S., J. Motiejūnaitė, I.S. Stepanchikova, D.E. Himelbrant & P. Czarnota 2012. New records of lichens and allied fungi from the Leningrad Region, Russia III. *Folia Cryptogamica Estonica* 49:31–37.
- Li, H.L. 1952. Floristic relationships between eastern Asia and eastern North America. *Transactions of the American Philosophical Society* 42: 371–429.
- Lobik, A.I. 1928. Materials for the mycological flora of the Kum River floodland according to study of 1925. In: *Materials on floristic and faunistic survey of the Tersky district*, pp. 13–29 (in Russian). [Лобик А.И. 1928. Материалы к микологической флоре плавень реки Кумы по обследованию 1925 года // Материалы по флористическому и фаунистическому обследованию Терского округа. С. 13–29].
- Lücking, R., B.P. Hodkinson & S. D. Leavitt 2016. The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota – approaching one thousand genera. *The Bryologist* 119(4): 361–416.
- Malençon, G. & R. Bertault 1969. Champignons du Maroc II. *Bulletin de la Société des Sciences Naturelles et Physiques du Maroc* 49(1–2):69–80.
- Malençon, G. & R. Bertault 1972. Champignons de la Péninsule Ibérique. IV. – Les Iles Baléares. *Acta Phytotaxonomica Barcinonensis* 11:1–64.
- Martius, H. 1817. *Prodromus florum Mosquensis*. Editio altera, Lipsiae: commercio industriae. 288 pp.
- Muchnik, E., L. Konoreva, S. Chesnokov, A. Paukov, A. Tsurykau & J. Gerasimova 2019. New and otherwise noteworthy records of lichenized and lichenicolous fungi from central European Russia. *Herzogia* 32:111–126.
- Notov, A.A., D.E. Himelbrant & G.P. Urbanavichus 2011. *The list of lichens and allied fungi of Tver Region*. Tver State University, Tver, 124 pp. (in Russian with English summary). [Нотов А.А., Гимельбрант Д.Е., Урбанавичус Г.П. 2011. Аннотированный список лишенофлоры Тверской области. Тверь: Тверской государственный университет. 124 с.].
- Popov, E.S. & S.V. Volobuev 2014. New data on wood-inhabiting macromycetes in key protected areas of the south-western part of the non-chernozem zone. *Mikologiya i fitopatologiya* 48(4):229–237 (in Russian with English summary). [Попов Е.С., Волобуев С.В. 2014. Новые данные о древообитающих макромицетах ключевых охраняемых природных территорий Юго-западного Нечерноземья // Микология и фитопатология. Т. 48, № 4. С. 229–237].
- Popov, E.S. 2012. New data on discomycetes of the Novgorod region. In: *Materials of the 2nd regional scientific-practical conference «Research and environmental activities in specially protected natural areas of the Novgorod Region», Valdai, 18–19 November 2011* (E.M. Litvinova, ed.), pp. 15–19, Velikiy Novgorod (in Russian). [Попов Е.С. 2012. Новые данные о дискомицетах Новгородской области // Исследования и природоохранные действия на особо охраняемых природных территориях Новгородской области: Материалы 2-й регион. науч.-практ. конф., г. Валдай, 18–19 ноября 2011 г. / Сост. и общ. ред. Е.М. Литвиновой. Великий Новгород. С. 15–19].
- Rodway, L. 1924. Tasmanian Discomycetes. *Proceedings of the Royal Society of Tasmania* 119:90–122.
- Rostrup, E. 1891. Tillaeg til "Grønlands svampe (1888)". In: *Conspectus florum groenlandicae. Pars tertia*, (J.S. Deichmann Branth, C. Grönlund, L. Kolderup Rosenvinge & E. Rostrup, eds.), pp. 593–643, I commission hos C.A. Reitzel, Kjøbenhavn.
- Sherwood, M.A. 1977. The ostropalean fungi. *Mycotaxon* 5: 1–277.
- Sirko, A.V. 1976. New and interesting species of ascomycetes in the flora of Urals. Discomycetes. *Novosti sistematiki nizshikh rastenii* 13:120–126 (in Russian). [Сирко А.В. 1976. Новые и интересные виды сумчатых грибов во флоре Урала. Дискомицеты // Новости систематики низших растений. Т. 13. С. 120–126].
- Smith, C.W., A. Aptroot, B.J. Coppins, A. Fletcher, O.L. Gilbert, P.W. James, & P.A. Wolseley 2009. *The lichen flora of Great Britain and Ireland*. British Lichen Society, London. 1046 p.
- Thümen, F. 1880. Beiträge zur Pilzflora Sibiriens. IV. *Bulletin de la Société Impériale des Naturalistes de Moscou* 55(2):198–233.
- Tranzschel, W. 1905. Materials for mycological flora of Russia. List of fungi collected in the Crimea. II. *Travaux du Musée Botanique de l'Académie Impériale des Sciences de St.-Peterbourg* 2:31–47 (in Russian). [Траншель В. 1905. Материалы для микологической флоры России. Список грибов, собранных в Крыму. II // Труды Ботанического Музея Императорской Академии Наук. Вып. 2. С. 31–47].
- Tranzschel, W. 1914. Die Pilze und Myxomyceten Kamtschatka's. In: *Expedition à Kamtchatka organisée par Th. P. Riabouchinsky avec le concours de la Société Impériale Russe de Géographie. Section de Botanique, vol. 2*, pp. 535–576, Moscow.
- Tranzschel, W. & I. Serebriankow 1912. *Mycotheca Rossica, sine Fungorum Rossiae et regionum confinium Asiae specimina exsiccata, fasc. 6–7*. Jaroslavl.
- Urbanavichene, I.N. 2011. First data on lichens of Zyuratkul national park (Chelyabinsk Region). *Novosti sistematiki nizshikh rastenii* 45:223–236 (in Russian with English summary). [Урбанавичене И.Н. 2011. Первые сведения о лишайниках национального парка «Зюраткуль» (Челябинская область) // Новости систематики низших растений. Т. 45. С. 223–236].
- Urbanavichene, I.N. & G.P. Urbanavichus 2014. Contribution to the lichen flora of the Achipse River valley (SW Caucasus, Krasnodarsky Krai). *Novosti sistematiki nizshikh rastenii*

- 48:315–326 (in Russian with English summary). [Урбанавичене И.Н., Урбанавичюс Г.П. 2014. К лихенофлоре долины реки Ачпсе (Юго-Западный Кавказ, Краснодарский край) // Новости систематики низших растений. Т. 48. С. 315–326].
- Urbanavichus, G.P. 2014. Additions to the lichen flora of Murmansk region. *Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii* 119(3):77 (in Russian). [Урбанавичюс Г.П. 2014. Дополнения к лихенофлоре Мурманской области // Бюллетень Московского общества испытателей природы. Отдел биологический. Т. 119, вып. 3. С. 77].
- Urbanavichus, G.P. & I.N. Urbanavichene 2015. A contribution to the lichen flora of Utrish Nature Reserve. *Turczaninovia* 18(2):86–95 (in Russian with English summary). [Урбанавичюс Г.П., Урбанавичене И.Н. 2015. Материалы к лихенофлоре заповедника «Утриш» // Turczaninovia. Т. 18, №2. С. 86–95].
- Urbanavichus, G.P. & I.N. Urbanavichene 2017. Contribution to the lichen flora of Erzi Nature Reserve, Republic of Ingushetia, North Caucasus, Russia. *Willdenowia* 47(3):227–237.
- Vaasma M., K.A. Kalamees & A. Raitviir 1986. *Macromycetes of the Caucasian State Reserve*. Valgus, Tallin, 106 pp. (in Russian). [Ваасма М., Каламеэс К. А., Райтвийр А. 1986. Макромикеты Кавказского государственного заповедника. Таллин: Валгус, 106 с.].
- Vasil'eva, L.N. 1939. Fungi of the Caucasian reserve. *Uchyonye zapiski kazanskogo gosudarstvennogo universiteta imeni V.I. Ul'yanova-Lenina* 99(1):1–66 (in Russian). [Васильева Л.Н. 1939. Грибы Кавказского заповедника // Учёные записки Казанского государственного университета им. В.И. Ульянова-Ленина. Т. 99, № 1. С. 1–66].
- Vasilyeva, L.N. & S.L. Stephenson 2010. Biogeographical patterns in pyrenomycetous fungi and their taxonomy. 1. The Grayan disjunction // *Mycotaxon* 114:281–303.
- Wedin, M., H. Döring & G. Gilenstam 2004. Saprotrophy and lichenization as options for the same fungal species on different substrata: environmental plasticity and fungal lifestyles in the *Stictis-Conotrema* complex. *The New Phytologist* 164:459–465.
- Wedin, M., H. Döring & G. Gilenstam 2006. *Stictis* s. lat. (Ostropales, Ascomycota) in northern Scandinavia, with a key and notes on morphological variation in relation to lifestyle. *Mycological research* 110:773–789.
- Wedin, M., H. Döring, K. Könberg & G. Gilenstam 2005. Generic delimitations in the family Stictidaceae (Ostropales, Ascomycota): the *Stictis-Conotrema* problem. *The Lichenologist* 37(1):67–75.
- Weinmann, J.A. 1828. Enumeratio Fungorum in agro Pawlowskiensi praecipue crescentium. *Sylloge plantarum novarum itemque minus cognitarum* 2:82–118.
- Wen, J. 1999. Evolution of eastern Asian and eastern North American disjunct pattern in flowering plants. *Annual Review of Ecology and Systematics* 30:421–455.
- Whitton, S.R., K.D. Hyde & E.H.C. McKenzie 1999. Microfungi on the Pandanaceae, a new species of *Stictis*. *Fungal Diversity* 2:169–174.
- Wu, Q. & G. M. Mueller 1997. Biogeographic relationships between the macrofungi of temperate eastern Asia and eastern North America. *Canadian Journal of Botany* 75(12): 2108–2116.
- Zervakis, G., P. Lizon, D. Dimou & E. Polemis 1999. Annotated check-list of the Greek macrofungi. II. Ascomycotina. *Mycotaxon* 72:487–506.