Freshwater centric diatoms from Middle Miocene deposits of the Khanka Depression, Primorye Territory (Far East of Russia)

Olesya Yu. Likhacheva¹, Aleksandra S. Avramenko^{2*},
Marina V. Usoltseva³ & Vladimir S. Pushkar¹

Olesya Yu. Likhacheva¹
e-mail: olesyalikh@gmail.com

Aleksandra S. Avramenko^{2*}
e-mail: alexa25rus@gmail.com

Marina V. Usoltseva³
e-mail: usmarina@inbox.ru

Vladimir S. Pushkar¹
e-mail: pushkar@fegi.ru

¹ Far East Geological Institute, FEB RAS,
Vladivostok, Russia

² Federal Scientific Center of the East
Asia Terrestrial Biodiversity, FEB RAS,
Vladivostok, Russia

³ Limnological Institute, SB RAS, Irkutsk,
Russia

* corresponding author

Manuscript received: 18.06.2021

Review completed: 29.10.2021

Accepted for publication: 12.11.2021

Published online: 15.11.2021

ABSTRACT

The centric diatom flora from the Middle Miocene deposits of the Khanka Depression of Primorye (Far East of Russia) was in detail studied. The centric diatoms were presented by 10 species and intraspecies taxa belonging to four genera (*Aulacoseira* Thwaites, *Alveolophora* Moiseeva et Nevretdinova, *Ellerbeckia* Crawford, *Melosira* Agard) and three orders (Aulacoseirales, Paraliales and Melosirales). A detailed study with the use of LM and SEM allowed us to establish the high morphologic variability of the predominant representatives of the genus *Aulacoseira*. The diagnoses for the *Alveolophora tscheremissinova*, *Ellerbeckia kochii*, *Melosira undulata* species were supplemented. The comparison of the taxonomic compositions of the centric diatoms from the Khanka complex with the even-aged complexes from other regions has shown the wide spreading of the genus *Aulacoseira* taxa in the freshwater bodies during this time period.

Keywords: diatom, Centrophyceae, *Aulacoseira*, *Alveolophora*, *Ellerbeckia*, *Melosira*, Middle Miocene, Khanka Depression, Khanka diatom flora, Primorye Region

РЕЗЮМЕ

Лихачева О.Ю., Авраменко А.С., Усольцева М.В., Пушкар В.С. Пресноводные центрические диатомеи из средне-миоценовых отложений Ханкайской депрессии Приморья (Дальний Восток России). Детально изучена центрическая диатомовая флора из среднемиоценовых отложений Ханкайской депрессии Приморья (Дальний Восток России). Центрические диатомеи представлены 10 видами и внутривидовыми таксонами, отнесенными к четырем родам: *Aulacoseira* Thwaites, *Alveolophora* Moiseeva et Nevretdinova, *Ellerbeckia* Crawford, *Melosira* Agard, трем порядкам Aulacoseirales, Paraliales и Melosirales. Детальное исследование с помощью СМ и СЭМ позволило установить высокую морфологическую вариабельность доминирующих представителей рода *Aulacoseira*. Дополнены диагнозы видов *Alveolophora tscheremissinova*, *Ellerbeckia kochii*, *Melosira undulata*. Сравнение таксономических составов центрических диатомей ханкайского комплекса с одновозрастными комплексами из других регионов показало широкое распространение таксонов рода *Aulacoseira* в пресных водоемах в этот период.

Ключевые слова: Диатомовые водоросли, Centrophyceae, *Aulacoseira*, *Alveolophora*, *Ellerbeckia*, *Melosira*, средний миоцен, Ханкайская депрессия, ханкайская диатомовая флора, Приморский край

The diatoms are one of the most informative groups for the paleoecological investigations (Stoermer & Smol 1999, Lewis et al. 2008, Haywood et al. 2009, Manoylov et al. 2009). These algae are sensitive indicators responding to the environmental changes (Fritz et al. 1991, 1993, Reid 2005). At this time, not only ecological structure of the paleocommunities of diatoms and number of valves in deposits but also morphological traits of phenotypes of individual taxa are taken into account (Douglas & Smol 1999, Cherepanova et al. 2007, Cherepanova et al. 2010). The examination of the Neogene diatoms (including the Middle Miocene ones) is necessary for determining the age and genesis of deposits and performing the paleogeographical reconstructions.

The Middle Miocene was marked by the series of the key climatic events which have transformed the global systems of the atmosphere and ocean circulations and resulted in the considerable changes in biota (Zachos et al. 2001,

Kürschner et al. 2008, Krapp & Jungclaus 2011). These events included the uplift of the Himalayas and Tibetan Plateau, change in intensity of the winter or summer Asian monsoon, and enhancing the aridity in Central Asia. The Himalayan Mountains began to form between 40 and 50 million years ago. The Himalayas had a height of more than 4,000 meters already by the end of the Early Miocene that led to formation of the regional climatic system on the outskirts of the Eurasian continent (Wang et al. 2003). The above mentioned events have considerably influenced on the climatic changes in the East Asia (Flower & Kennett 1994, Zhisheng 2000, Zhisheng et al. 2001, Jiang & Ding 2008, Royden et al. 2008). The Middle Miocene Climatic Optimum (14.5–17 Ma, maximum – 16 Ma) is one of the most important events and also the warmest period in the past 30 million years (Kennett 1982, Tsuchi & Ingle 1992, Zachos et al. 2001, Shevenell & Kennett 2004). The tempe-

tures of the sea water and air over the land surface were higher from the Eocene to the present day (Tanai & Huzio-ka 1967, Kennett 1982, Tsuchi & Ingle 1992, Schoell et al. 1994). The similar trends in the climate changes were fixed for many regions of the Earth including Primorye Territory (Pavlyutkin 2005, Pavlyutkin et al. 2004). At the same time, the intensification of the East-Asian monsoon and emergence of the systems of warm and cold currents in the Sea of Japan have reduced the influence of the Middle Miocene Climatic Optimum in the south of the Russian Far East (Likhacheva et al. 2009).

On the territory of Russia, the Middle-Miocene continental deposits containing the diatom algae were found in several areas. For the first time, they were investigated in 1952 by A.P. Jousé (1952) from the locations in the vicinity of Khanka Lake (Primorsky Region). Jousé has promoted the development of studying the Middle Miocene diatom taxa by describing and documenting new taxa and by illustrating a number of previously undescribed forms. She has established the diatom complexes characteristic of the Middle Miocene. The stratigraphy of the USSR has entered a new stage of development due to study of the diatom complexes and its extensive correlation.

In the 1960s, there was a significant increase in the scope of geological surveys on the territories occupied by the continental Neogene deposits including Miocene ones: the south of the Khabarovsk and the Primorsky Territories; Tunka Basin, Transbaikalia in Baikal Region; Irtys River, Tym, West Siberian Plain in the Western Siberia (Proshkina-Lavrenko 1974) and Yamal Peninsula (Polyakova & Danilov 1989). By the 1970s, a number of deposit sites were analyzed for diatoms and the first diatom Neogene biostratigraphy in the USSR has appeared. The reconstruction of Neogene climate was also based upon diatom analyses. The remains of centric diatoms in these sediments are one of the major organogenic components. The studies of Neogene diatoms in the USSR were in detail described in the extensive report published in 1974 (Proshkina-Lavrenko 1974).

A new stage in research of the continental deposits from East of Russia is associated with the beginning of the 21st century. The Middle-Late Miocene continental sediments were studied in Transbaikalia: Vitim Upland (Chernyayeva et al. 2007, Rasskazov et al. 2007, Usoltseva et al. 2010, Usoltseva & Khursevich 2013), Chara Basin (Chernyayeva & Moiseeva 2003, Enikeev 2008), Tunka Basin (Proshkina-Lavrenko 1974, Popova et al. 1989, Likhoshway et al. 1997), Barguzin Basin (Hassan et al. 2019, Usoltseva et al. 2020) and the Southern Baikal hollow (Chernyayeva & Popova 1993, Mats 2013, Rasskazov et al. 2014) and Upper Miocene deposits from Baikal Lake (Khursevich et al. 2005, Kuzmin et al. 2009).

The freshwater diatom flora was also found in the Lower Miocene deposits from Yamato and Krishtovich Rises in the Sea of Japan (Usoltseva & Tsoy 2010, Tsoy & Usoltseva 2015, Tsoy 2017). New fossil diatom species of *Aulacoseira* (Tanaka et al. 2008, Tanaka & Nagumo 2011a, b), *Actinocyclus* (Tanaka & Nagumo 2014, Hayashi et al. 2012) and *Tetracyclus* (Saito-Kato et al. 2011) from Miocene deposits of Japan and Sea of Japan were described.

Diatoms were also found in the Middle Miocene Shanwang Formation, Shandong Province, East China (Li et al. 2010).

The studies of the Middle Miocene diatom flora of the Khanka Lake area begun by Jousé (1952) were continued by Moiseeva (Moiseeva 1971, 1995). Later, Tsarko (Moiseeva 1971, 1995, Moiseeva & Tsarko 1990), Pushkar, Cherepanova and Likhacheva (Pavlyutkin et al. 1993, Pavlyutkin et al. 2004, Pushkar & Likhacheva 2007) have also joined with the research. The valve morphology of the genus *Tetracyclus* Ralfs from these deposits was studied by Usoltseva and Dubrovina with co-authors (Dubrovina et al. 2011, Dubrovina et al. 2014). However, the dominant *Aulacoseira* Thwaites and other centric taxa were not previously described. The appropriate morphological characters for taxonomic separation of *Aulacoseira* are still largely unclear. The species of the genus *Aulacoseira* are quite variable. Plasticity of morphological features makes identification of *Aulacoseira* species difficult. The studies carried out for ultrastructure of "prae" of *Aulacoseira* taxa using the electron microscopy are few in number (Usoltseva & Likhoshway 2005, 2007, Usoltseva et al. 2011). The accurate identification of diatom species is necessary for taxonomy, stratigraphy, paleoenvironmental reconstructions and decipher of the past climatic changes. The electronic microscopy can partially resolve these problems providing the additional information for classification of *Aulacoseira*.

The study purpose is a detail study of the Middle Miocene freshwater centric diatoms from the Khanka deposits (Primorye Territory, Far East of Russia) formed at temperatures higher than today using both light and electron scanning microscopes.

STUDY AREA

The Khanka Depression is situated in the west of Primorsky Territory (Fig. 1). The deposits forming the hollow have resulted from the tectonic and magmatic processes and presented by the Mesozoic and Cenozoic sedimentary and volcanogenic rocks. Of the Late Cretaceous–Early Paleogene, the tectonic-magmatic activation is characteristic. In the Eocene-Oligocene, the depression plunges, alluvial and lacustrine sediments accumulate in it. A volcanism with the felsic and moderately felsic eruptions was one of the most important events in the Miocene. From the beginning of the Middle Miocene, the process of sedimentation in lakes located in the floodplain of a large river or in its delta continues. In the Pliocene, acidic volcanism gives way to basaltic volcanism. The depression plunging intensity increases causing the river system restructuring. In the Pliocene, a geological and geomorphological situation similar to the present one has developed. However, the formation of some morphostructures has continued in the Quaternary as well. This is shown by volcanism, seismic activity which caused the changes in the area of Khanka Lake.

The modern terrain of the Khanka Depression is low-contrast and characterized by low-mountain ridges with flattened watersheds and gentle slopes. Absolute relief elevations of the watershed areas within the limits of depression are 120–140 m, while 400–600 m in its framing. The maximum elevation is 780 m a.s.l. The climate is monsoonal. Win-

ters are dry with a lack of snowfall; summers are warm and rainy. The average annual temperature is $+3^{\circ}\text{C}$, the average temperature of the coldest month (January) is -18.4°C , and the warmest month is August ($+20.7^{\circ}\text{C}$). The average annual precipitation is 527 mm of which 40 mm falls in winter.

MATERIAL AND METHODS

The investigated samples were collected from the Middle Miocene diatomite-bearing deposits in the Khanka Depression (Primorye Territory, Far East of Russia). The five samples were taken by Pavlyutkin in 2004 from section 9149 (diatom collection of Prof. Pushkar) and thirty five samples were taken by Likhacheva in three neighboring sections (Fig. 1): № L-03/2 ($45^{\circ}09'14.3''$ and $132^{\circ}00'23.3''$) and L-04/1 ($45^{\circ}09'11.8''$ and $132^{\circ}00'126.00''$) in 2008, L-02 ($45^{\circ}01'00.0''$ and $132^{\circ}00'25.1''$) in 2010–2011. The sections are located on the western shore of Khanka Lake between the settlements of Turiy Rog and Novokachalinsk (north-west of the Khanka Depression). According to Pavlyutkin, this flora is named to as Khanka flora (2005).

The western shore of Khanka Lake in segment being of interest to us is the abrasion scarp with a narrow broken strip of sand and pebble beach (Fig. 1). The deposits are represented by pebbles, sandstones and thin-layered tuff-alevrolites and tuffaceous diatomites. This sedimentary stratum belongs to the Novokachalinskaya Formation (Pavlyutkin et al. 2004). The deposits of this suite overlie the Upper Pliocene coal-bearing deposits and are overlapped by the Upper Pliocene clays. The age of the Novokachalinskaya Formation rocks was determined according to data of

the paleobotanic, palynological and diatom analyses as the Middle Miocene (Pavlyutkin et al. 2004, Pavlyutkin 2005). The Middle Miocene volcanites from Novokachalinskaya Formation were dated 18.1–14.9 Ma (Pavlyutkin et al. 1993, Likhacheva et al. 2009). The Middle Miocene corresponds to the Langhian and Serravallian stages according to the International Chronostratigraphic Scale (Cohen et al. 2021).

The samples were treated by the standard procedure using hydrogen peroxide (Proshkina-Lavrenko 1974). Taxonomic analysis, counting of diatom valves, and measurements of the diameter, length, and width of the valves were performed on Amplival Carl-Zeiss and Axioskop 40 Carl Zeiss light microscopes (LM) using a 18×18 mm cover glass at a magnification of $\times 1000$ with an immersion liquid at the Center for Collective Use of Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far East Branch, Russian Academy of Sciences. 250–300 diatom valves were counted to determine the percentage of individual species in diatom taphocenoses (diatom fossil community). Morphological features of valves were also examined with a Carl Zeiss EVO 40, Merlin and Philips 525M scanning electron microscope (SEM) at a magnification of up to 15,000. The sediment remaining after the preparation of the specimen for LM studies was dried. The resulting powder was applied on the special tables with glued bilateral carbon tape. The samples were sputter coated with gold in a vacuum chamber (JEOL JFC-1600 auto fine coater).

The taxonomical features of the genus *Aulacoseira* are form and dimensions of the valves, a form of rimoportulae and their location on the valve as well as the frequency of particular valve characters such as the number of rows of areolae in $10\text{ }\mu\text{m}$ and the areolar density in $100\text{ }\mu\text{m}^2$ on the mantle and ratio between valve height and diameter (Krammer & Lange-Bertalot 1991). We have also measured a height and diameter of valves and determined the correlation coefficients (Cv) of these features (Ershova 2015).

Classification of centric diatoms follows the systematics as proposed by Round et al. (1990) was modified by Nikolaev & Harwood (2001). All described species are recognized taxonomically valid (Guiry & Guiry 2021).

RESULTS

Diatoms of the Coscinodiscophyceae class dominate in the Middle Miocene deposits of the Khanka Depression. This class is represented by two subclasses, three orders and four genera, which include 10 species and intraspecific taxa.

The Paraliophycidae subclass is represented by two orders (Aulacoseirales and Paraliales); each of them contains one family (Aulacoseiraceae и Radialiplicataceae). The Aulacoseiraceae includes two genera: *Aulacoseira* and *Alveolophora* Moiseeva et Nevretdinova, while the Radialiplicataceae one genus (*Ellerbeckia* Crawford). One species, three varieties and six forms were identified in the genus *Aulacoseira*:

- A. praegr anulata* var. *praeislandica* f. *praeislandica* (Jousé) Moiseeva
- A. p.* var. *praeislandica* f. *curvata* (Jousé) Moiseeva
- A. p.* var. *praegr anulata* f. *praegr anulata* (Jousé) Simonsen
- A. p.* var. *praegr anulata* f. *curvata* (Jousé) Simonsen
- A. p.* var. *praeangustissima* f. *praeangustissima* (Jousé) Moiseeva
- A. p.* var. *praeangustissima* f. *curvata* (Jousé) Moiseeva.

Figure 1 Study area. Locations of sampling sites are shown with black dots on the western shore of the Khanka Lake of Primorye

Two species were identified in *Alveolophora*:

A. tscheremissinovi Khursevich
A. khursevichiae Usoltseva, Pushkar & Likhacheva.

The last mentioned species was recently described and found only in the Middle Miocene deposits of the Khanka Lake area. One species was identified in *Ellerbeckia*:

E. kochii (Pantocsek) Lupikina.

The subclass Melosirophyceae is represented by one order containing one family (Melosiraceae) represented by the genus *Melosira* Agardh, which includes only one species (*Melosira undulata* (Ehrenberg) Kützing).

This flora may be treated as the *Aulacoseira* flora because the rock-forming taxa are represented by the genus *Aulacoseira*. The dominant taxon is *A. praegratulata* var. *praeislandica* f. *praeislandica*. Its percentage in the diatom paleocommunities reaches 84.7 %. *A. praegratulata* var. *praeislandica* f. *curvata* has percentage of up to 7.27 %, *A. praegratulata* var. *praegratulata* f. *praegratulata* up to 2 %. Other taxa occur in low frequencies (<1%).

SYSTEMATIC DESCRIPTIONS

AULACOSEIRACEAE

Aulacoseira praegratulata (Jousé) Simonsen, 1979, Bacillaria 2: 62.

Description. Frustules are short or long cylinders. The diatom can be found from short chains of 2–3 cells to long chains of 4–15 cells. Valves are thick-walled. They are straight or curved on central axis. The areolae on the mantle are situated in straight or curved rows. The areolae are varying from rounded to round-square in shape. Detail on the external valve surface showing the areolae occluded by rotulae (in the majority of valves the membrane is destroyed). Sometimes on a mantle are available small perforations, randomly located between rows of areolae. Structureless collar is quite short. The ring furrow usually isn't expressed. Rimoportulae are located along one ringleist. The spines occur regularly between the last areola in a row on the mantle. Spines are tapering or have a spade-shape at the end.

It's a freshwater extinct species, known from Oligocene and Early Pleistocene sediments (33.9–0.781 Ma).

var. *praegratulata* (Jousé) Simonsen

Frustules are long cylinders. Valves are curved or straight on central axis, 5.73–10 µm in diameter. Areolae on the flat valve face are situated distantly. Valves are 5.97–21.88 µm in

height. Rows of mantle areolae are straight or curved to the right, 9–14 rows in 10 µm, the areolar density in 100 µm² are 76–180. The ratio between valve height and diameter varies from 1.3 to 3.27 (average 2.37). The collar is mostly short. Spines are spade-shaped.

f. *praegratulata* – *Melosira prae-gratulata* [(Ehrenberg) Ralfs] Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241. – *M. praegratulata* Jousé f. *praegratulata*, 1971, in Moiseeva: 26–27

Frustules along central axis are straight (Fig. 2: A–D; Table 1).

f. *curvata* – *Melosira prae-gratulata* [(Ehrenberg) Ralfs] f. *curvata* Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241

Frustules along central axis are curved (Fig. 2: E).

var. *praeislandica* (Jousé) Moiseeva, in Glezer et al. 1992: 85

Frustules are short or long cylinders, in valve view are round. Valves are thick-walled. They are curved or straight on pervalvar axis, 4.84–27 µm in diameter (diameter is equal to height of a valve or exceeds it). Valves are 2.0–17.95 µm in height. Rows of mantle areolae are straight or curved to the right, 8–12 rows, sometimes to 18, in 10 µm, the areolar density in 100 µm² are 52–208, sometimes to 236. Areolae on the flat valve face are situated distantly or closely. The ratio between valve height and diameter varies from 0.14 to 2.67 (average 0.87). The collar is short.

f. *praeislandica* – *Melosira prae-islandica* (O.Müller) Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241. – *M. prae-distans* [(Ehrenberg) Kütz.] Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241. – *M. praeislandica* Jousé f. *praeislandica* in Moiseeva, 1971: 28–29. – *M. praedistans* f. *praedistans* Moiseeva, 1971: 25

Frustules along central axis are straight (Fig. 3: A–C, G–J, K (top), L–N, Q–T; Table 1).

f. *curvata* – *M. prae-islandica* f. *curvata* Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241. – *M. praedistans* f. *curvata* Moiseeva, 1971: 26

Frustules along central axis are curved (Fig. 3: D–F, K (bottom)).

var. *praeangustissima* (Jousé) Moiseeva, in Glezer et al. 1992: 84

Frustules are long and narrow cylinders. Valves are curved or straight on central axis, 2.92–7.36 µm in diameter. Valves are 14.04–23.5 µm in height. Valves are much longer,

Table 1. Observed morphological variation in the varieties of the species *Aulacoseira praegratulata*.

Taxa	Valve mantle height (h)				Valve diameter (d)				Height / Diameter ratio (middle)	Areolar density in 100 µm ² on the mantle	Number of rows in areolae in 10 µm
	Min–Max middle, µm	Number of valve measured	σ	Cv	Min–Max, middle, µm	Number of valve measured	σ	Cv			
var. <i>praegratulata</i> f. <i>praegratulata</i>	5.97–21.88 16.6±0.6	37	3.5	21.3	5.73–10.0 7.4±0.2	34	1.0	14.1	1.3–3.27 (2.38)	76–180	9–14
var. <i>praeislandica</i> f. <i>praeislandica</i>	1.86–17.95, 10.1±0.1	406	2.9	28.3	4.84–27.0, 13.3±0.2	478	3.6	26.8	0.14–2/67 (0.87)	52–208 (236)	8–12 (18)
var. <i>praeangustissima</i> f. <i>praeangustissima</i>	14.04–23.5, 18.8±0.8	12	2.8	14.9	2.92–7.36, 5.14±0.2	17	1.0	18.0	2.84–6.34 (4.14)	36–112	9–10

Note. Cv is the coefficient of variation; σ is the standard deviation; ± the standard error of the mean

Figure 2 *Aulacoseira praegranulata* var. *praegranulata* f. *praegranulata* (Jousé) Simonsen. A–D – valves are straight along central axis; A – valve mantle view (LM); B, D – valve mantle view, the areolae on the mantle are arranged in curved rows (SEM); B – perforations irregularly located between the rows of areolae on the valve mantle; C – valve with short structureless collar (SEM); B–D – valves with spade-shaped linking spines (SEM). *Aulacoseira praegranulata* var. *praegranulata* f. *curvata* (Jousé) Simonsen. E – valve is curve along central axis (LM)

then wide. Rows of mantle areolae are straight or curved to the right, 9–10 rows in 10 µm, the areolar density in 100 µm² are 36–112. Areolae on the flat valve face are situated distantly. The ratio between valve height and diameter varies from 2.84 to 6.34 (average 4.14). The collar is short.

f. *praeangustissima* – *M. prae-granulata* var. *praeangustissima* [(O.Müller) Hust.] Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241

Frustules along central axis are straight. (Fig. 4: A, B, D; Table 1).

f. *curvata* – *M. prae-granulata* var. *praeangustissima* [(O.Müller) Hust.] f. *curvata* Jousé, 1952, Trudy instituta geografii AN SSSR 6(51): 241

Frustules along central axis are curved. (Fig. 4: C).

Comment. Auxospores were found. Auxospores have hemispherical front part of valve and differ from valve more rough structure and well expressed ring-shaped diaphragm.

Alveolophora tscheremissinovae Khursevich. This name is currently regarded as a synonym of *Miosira tscheremissinovae* (Khursevich) Khursevich, 2012, in Khursevich & Kociolek 141: 347

Description. Frustules are short cylinders. Frustules in valve view are round, 8.4–11.3 µm in diameter. Valves are thick-walled. Valves are 1.5–2.9 µm in height. Mild areolae on the valve face are situated in the near-margin surface, in radial rows, 8 rows in 10 µm. Radial rows of areolae are located in narrow (3–4 rows) sectors, which are divided by direct costal thickenings. Rimoportulae have the form of curved tube (3–5 on a valve), form ring in margin (if 5) or central (if 3) zone of a face part of valves. At margin zone of the valve there are poorly expressed pseudosepta. Areolae on the mantle are situated in straight rows (14–18 in 10 µm, two areolae in each row). Spines are short, spade-shaped are located on border of face part of valves and mantle (Fig. 5).

Freshwater extinct species and known from Miocene deposits (23.03–5.33 Ma).

Comment. The smallest valves we measured were 8.4 µm in diameter. It is smaller than that of the typical forms described in Khursevich (1994). The valves have 3–5 rimoportulae, which is less than of typical forms. The valves have 14–18 areolae on the mantle in 10 µm, two areolae in each row. The typical forms have 8–12 areolae on the mantle in 10 µm, 2–3 areolae in each row.

Alveolophora khursevichiae Usoltseva, Pushkar & Likhacheva (Usoltseva et al. 2018, Fig. 2–24).

RADIALIPLICATACEAE

Ellerbeckia kochii (Pantocsek) Lupikina 1991: 59. – *Melosira kochii* Pantocsek, 1905: 62. – *M. arenaria* var. *hungarica* Pantocsek, 1905: 59. – *M. varennarum* Peragallo et Hérilbaud, in Hérilbaud, 1893: 189. – *M. scabrosa* Østrup sensu Moiseeva, 1971: 30–31. – *Paralia varennarum* (Peragallo et Hérilbaud) Loseva, 1981: 77. – *P. kochii* (Pantocsek) Moiseeva, 1987, in Moiseeva & Genkal: 1502

Description. Frustules tightly united by linking structures on valve faces to form long chains. Frustules are broadly cylindrical, complex. Valves are 15.8–57.1 µm in diameter, 4–17 µm in height. Linking structures or relief valve represented by radial ridges of different lengths (9–12 in 10 µm) in the submarginal zone or the valve face and corresponding structures of intaglio valve are radial grooves. Ridges are up to 1/3 or 1/2 of the valve radius. Remainder of the valve face is covered by unevenly located tuberos, different outlines silicified thickenings. Valve mantle penetrated by pores 17–30 in 10 µm of the vertical row. Vertical rows of pores (20–25 in 10 µm) separated by narrow ribs. Externally, relief valve with stepped mantle concave, flat are more often, intaglio valve with plain mantle convex. Original tube processes (2–3 in 10 µm) formed by six ribs curved spirally upwards and alternate with pores on internal surface or valve mantle. Externally tube processes terminate

Figure 3 *Aulacoseira praeislandica* var. *praeislandica* f. *praeislandica* (Jousé) Moiseeva (A–C, G–T, K–N, Q–T, V): valves are straight along central axis; A, C, F, L, M, R, U, T, V – valve mantle view, the areolae on the mantle are arranged in straight rows (A, L, V – LM; C, F, M, R, U, T – SEM); B, G, N – valve mantle view, the areolae on the mantle are arranged in curve rows (B – LM; G, N – SEM); L–N, Q, V – low-cylindrical frustules (N, Q – SEM; A–G, L–N, R, V – chains consisting of some valves; C, F, Q – perforations irregularly located between the rows of areolae on the valve mantle; K (valve located at the photo top) – ringleist of straight valve (SEM); H–J – external valve face (H – LM; I–J – SEM); O – post-auxospore chain with hemispherical initial valve (SEM); P – spore (SEM); S, T – fragments showing views of broken ringleist with rimoportulae (arrows) (SEM); U – valve mantle view with the external opening of the rimoportula (arrow) (SEM); Q, R – valves with tapering linking spines; C, F, G, M, N, U – valves with spade-shaped linking spines; *Aulacoseira praeislandica* var. *praeislandica* f. *curvata* (Jousé) Moiseeva (D–F, K): valves are curve along central axis; D–F – valve mantle view (D – LM; E–F – SEM), K (valve located at the photo bottom) – ringleist of curve valve (SEM).

by openings. They are located irregularly in two rings. Mantle edge is crenulate. Crenulations entwine with similar structures on valvocopula. Valvocopula ornamented of thin pores: 12–25 pores in 10 µm or vertical rows, 16–30 pores in 10 µm or horizontal rows. Valves united by means or a

circular interlocking diaphragm, located along the mantle rim in valves plane. Diaphragm relief represented by a system or alternating shallow radial furrows and ribs (Fig. 6). Freshwater extinct species and known from Miocene, Pliocene (5.332–1.806 Ma), Lower Pleistocene (1.806–0.781 Ma).

Figure 4 *Aulacoseira praeangustissima* f. *praeangustissima* (Jousé) Moiseeva (A, B, D): valves are straight along central axis; A, B, D – long cylindrical frustules (A, B – LM; D – SEM); D – valve with high collar and perforations irregularly located between the rows of areolas on the valve mantle; *Aulacoseira praeangustissima* f. *curvata* (Jousé) Moiseeva (C): valve is curve along central axis; C – valve with spade-shaped linking spines (LM)

Comment. The smallest valves we measured were 15.8 µm in diameter and 4 µm in height. It is smaller than that of the typical forms: 30–70 µm in diameter, 10–17 µm in height (Kuzmin et al. 2009), 34–65 µm in diameter, 15–17 µm in height (Moiseeva 1971).

MELOSIRACEAE

Melosira undulata (Ehrenberg) Kützing, 1844: 54. – *Gallionella undulata* Ehrenberg, 1840: 211.

Description. Frustules are barrel-shaped, with a flat valve face forming a very small pseudosulcus. Valves 14.35–54.1 µm in diameter, ornamented with radiating rows of striae of different lengths on the face, about 12–16 in 10 µm. Striae on the valve face are dichotomously branched and radiate from a hyaline central area. Valves are 7.65–31.28 µm in height. The mantle ornamentation is composed of parallel fine punctae arranged along the pervalar axis, 16–22 in 10 µm. Rimoportulae form a ring one-fourth to one-third the distance between the valve and the girdle edge. Several rimoportulae are also present on the surface of the valve. Although the external mantle walls are parallel, the inner surface of the wall is clearly undulated. The undulation is seen as a varying thickness of the wall, with the thickest portion about two-thirds below the valve face. Spines are small and barely distinguishable (Fig. 7). Freshwater species and known from Miocene to the present.

Comment. The smallest valves we measured were 7.65 µm in height. It is smaller than that of the typical forms: 20–35 µm (Hustedt 1930), 18–32 µm (Moiseeva 1971), 28–47 µm (Krammer & Lange-Bertalot 1991).

DISCUSSION

The researchers studied previously the Khanka diatom flora have also noted a dominance of centric taxa in the Middle Miocene deposits. Jousé (1952) has found 12 species

and intraspecific taxa of the genus *Melosira* (Table 2). Moiseeva (1971) has identified eight taxa belonging to this genus (Table 2). Pushkar and Cherepanova have studied the Middle Miocene deposits from natural rock outcrops and sediment cores. The taxonomic composition of the centric flora was supplemented. Pushkar and Cherepanova have identified 14 taxa of the *Melosira*, *Aulacoseira*, *Paralia*, *Alveolophora* (Pavlyutkin et al. 2004). Likhacheva et al. (2009) used the LM and SEM to determine the taxonomic composition of the Khanka flora and found 15 centric taxa relating to genera *Melosira*, *Aulacoseira*, *Ellerbekia*, *Alveolophora*. Likhacheva and colleagues (Likhacheva et al. 2009, Likhacheva 2013) have provided the Neogene diatom biostratigraphic scheme for Primorye, including the Middle Miocene of the Khanka Depression, using the latest evidences in taxonomy and biostratigraphic distribution of Neogene diatoms. According to the data presented above, the Khanka centric diatom flora has shown low taxonomic diversity: 10 diatom species and intraspecific taxa belonging to four genera (Table 2).

A special attention in this study was paid to investigating the dominant representatives of the genus *Aulacoseira* belonging to the “prae” group of which a great number of intermediate forms is characteristic. The species of *Aulacoseira* described in this report have been separated by more or less obvious, even if variable, morphological discrepancies in height and diameter of valves (Table 1). Analysis of the valve characteristics for three *Aulacoseira* varieties showed that diameter of *A. praeangustissima* f. *praeangustissima* is less variable (Cv=14.1) than the height of valve (Cv=21.3). Both diameter (Cv=18.0) and height of valve (Cv=14.9) of *A. praeangustissima* f. *praeangustissima* are constant. For *A. praeangustissima* f. *praeislandica*, both diameter (Cv=26.8) and height of valve (Cv=28.3) are variable. If the value of the variation coefficient does not exceed 33 %, the sample is

Figure 5 *Alveolophora tscheremissinovae* Khursevich: A, G – valve mantle view (A – LM, G – SEM), B–F, H – internal valve views with rimoportulae (arrows) (SEM); B–D, G–H – valve with short spatulate linking spines, H – chain consisting of some valves

homogeneous (Ershova 2015). The coefficients of variation for the valve diameter and height of *A. praegr anulata* var. *praeislandica* f. *praeislandica* are less than 33 %, but higher than those for other two varieties. This value of the coefficient is determined by a wide range of valve mantle and diameter; this taxon has a high mantle and small diameter (Fig. 3A–D, G) and low mantle and large diameter (Fig. 3L–N, V, Q). Jousé (1952) has also found the diatoms with similar characteristics in the Miocene sediments of the Khanka Lake area. Jousé has described two species: *Melosira prae-islandica* (O. Müller) Jousé and *Melosira prae-distans* Jousé (hereinafter the spellings of taxa are according to the authors).

A large number of representatives of the genus *Melosira* was transferred by Simonsen (1979) to the genus *Anuloseira*.

Thus, this genus included only *A. praegr anulata* (Jousé) Simonsen with form of *curvata* and *A. praeislandica* (Jousé) Simonsen. Temniskova & Vodenicharov (1981) have investigated *A. praeislandica* and *Melosira praedistans* in the Lower Pliocene freshwater diatomite from South-Western Bulgaria. The ultrastructures of valves and biometric investigations did not show the clear differences between the species. The authors concluded that these taxa belong not only to the same genus but also to the same species of *A. praeislandica*. As a result, the diagnosis of this species was expanded. Officially, these two species were later included in synonymy of *A. praegr anulata* var. *praeislandica* f. *praeislandica* (Glezer et al. 1992). Usoltseva & Likhoshvay (2005) have studied the original material from the Middle Miocene deposits of the Khanka Depression

Figure 6 *Ellerbeckia kochii* (Pantocsek) Lupikina: A, B, E – valve face views with short radial ridges (A – LM; B, E – SEM); B, E, G – valvo-copula consists of thin pores in vertical rows (G – SEM); B, E – radial ridges in the valve submarginal zone (SEM); C, D, F – structure of tube processes on internal valve mantle (arrows) (SEM); E – valve mantle view with the external opening of tube processes (arrows); C, D – internal valve mantle with two rings of tube processes (arrows); D, F – internal valve mantle structure

taken by Jousé in 1952. They have investigated all species of *Melosira praedistans*, *M. praeislandica* and *M. praegr anulata* by SEM. It was found that there are no clear differences between these species in both the valve forms and the elements of the frustules ultrastructure (Usoltseva & Likhoshway 2005). The studied diatom taphocenoses are a set of valves which were accumulated in sediments over many thousands and even millions of years. They include valves formed in different seasons of year and at different stages of the life cycle of algae, thus, probably, we observe a wide range of fluctuations in the height and diameter of the valves.

One of the criteria for the identification of *Aulacoseira* is the valve height to valve diameter ratio (H/D) (Table 1). The H/D of *A. praegr anulata* var. *praegr anulata* f. *praegr anulata* is 1.3–3.27, the average value of H/D ratio is 2.37, and for

most valves this indicator is within 2.21–2.95. The H/D of *A. praegr anulata* var. *praeangustissima* f. *praeangustissima* is 2.84–6.34, in average, 4.14 and most valves are limited to 3.23–4.49. The H/D of *A. praegr anulata* var. *praeislandica* f. *praeislandica* is 0.14–2.67, in average, 0.87.

The form and localization of rimoportulae are important diagnostic features of *Aulacoseira* (Likhoshway & Crawford 2001). The investigation showed that the rimoportulae of the studied species are located on the border with the ringleist (Fig. 3S). The similar "sessile" rimoportulae were noted (Usoltseva & Likhoshway 2005) for the *Aulacoseira* species from the Jousé original material.

The diatoms have shortish structureless collar (Fig. 2B–D, Fig. 3C,F,G,U, Fig. 4A), the sulcus was not usually obscure; the areoles were locular (Fig. 2, 3, 4).

Figure 7 *Melosira undulata* (Ehrenberg) Kützing: A, C–E – external valve face (A, C–D – LM; E – SEM); B, F, G – valve mantle view showing undulate inner surface of the wall (LM); D – external valve face views showing the position of rimoportulae

The diagnoses of *Alveolophora tscheremissinovae*, *Ellerbeckia kochii* and *Melosira undulata* were supplemented. It has been established that the minimum valve diameter of *A. tscheremissinovae* (8.4 µm) is smaller than that of the typical forms – 10–24 µm (Khursevich 1994). The valve had a smaller number of rimoportulae (3–5), whereas the typical forms had 4–12. On the contrary, the number of areola rows on the mantle is larger (14–18 rows per 10 µm), while 8–12 rows in 10 µm in the typical forms. The valves of *E. kochii* had a smaller diameter (from 15.8 µm) than that indicated in the diagnosis (from 30 µm) (Glezer et al. 1992)

and a lower height of valve (from 4 µm to 17 µm). The height of *E. kochii* valve is 10–15 µm (Glezer et al. 1992).

According to the diagnoses of various researchers, the minimum height of *M. undulata* valve is 20 µm (Hustedt 1930), 18 µm (Moiseeva 1971) and 28 µm (Krammer & Lange-Bertalot 1991). *M. undulata* from Novokachalinskaya Formation had a lower valve height which is 7.65 µm.

The Khanka diatom flora has been studied for more than 70 years, but there are still many questions concerned its systematics and taxonomic composition. First of all, the western lake shore exposed the Middle Miocene deposits

Table 2. Freshwater centric diatom flora from Middle Miocene deposits of the Khanka depression based on this paper and references

Centric taxa	Present study	Jousé (1952)	Moiseeva (1971)	Pushkar, Cherepanova (Pavlyutkin et al. 2004)	Likhacheva (2013), Likhacheva et al. (2009)
<i>Aulacoseira praegranulata</i> var. <i>praeislandica</i> f. <i>praeislandica</i>	+	* <i>Melosira prae-islandica</i> , <i>M. prae-distans</i>	* <i>M. praeislandica</i> f. <i>prae-islandica</i> , <i>M. praedistans</i> f. <i>praedistans</i>	<i>A. praegranulata</i> var. <i>praeislandica</i> f. <i>praeislandica</i> , <i>A. praedistans</i>	+
<i>A. praegranulata</i> var. <i>praeislandica</i> f. <i>curvata</i>	+	* <i>M. prae-islandica</i> f. <i>curvata</i> Jousé	* <i>M. praedistans</i> f. <i>curvata</i> , * <i>M. praeislandica</i> f. <i>curvata</i>	+	+
<i>A. praegranulata</i> var. <i>praegranulata</i> f. <i>praegranulata</i>	+	* <i>M. prae-granulata</i>	* <i>M. praegranulata</i> f. <i>prae-granulata</i>	+	+
<i>A. praegranulata</i> var. <i>praegranulata</i> f. <i>curvata</i>	+	* <i>M. prae-granulata</i> f. <i>curvata</i>	* <i>M. praegranulata</i> f. <i>curvata</i>	+	+
<i>A. praegranulata</i> var. <i>praeangustissima</i> f. <i>praeangustissima</i>	+	* <i>M. prae-granulata</i> var. <i>prae-angustissima</i>	—	+	+
<i>A. praegranulata</i> var. <i>praeangustissima</i> f. <i>curvata</i>	+	* <i>M. prae-granulata</i> var. <i>prae-angustissima</i> f. <i>curvata</i>	—	+	—
<i>A. moiseevii</i> Akiba et Tsoy	—	—	—	+	—
<i>A. italica</i> var. <i>italica</i> (Kützing) Simonsen	—	—	—	+	+
<i>A. italica</i> var. <i>italica</i> f. <i>curvata</i> (Pantocsek) Davydova	—	—	—	—	+
<i>Ellerbeckia kochii</i>	+	—	<i>Melosira scabrosa</i>	<i>Paralia kochii</i>	+
<i>Ellerbeckia arenaria</i> var. <i>teres</i>	—	—	—	+	+
<i>Melosira undulata</i>	+	+	<i>Melosira undulata</i> var. <i>undulata</i> (Ehrenberg) Kützing	+	<i>Melosira undulata</i> var. <i>undulata</i>
<i>Melosira italica</i> f. <i>miocenica</i>	—	+	—	—	—
<i>M. italica</i> var. <i>tenuissima</i> f. <i>miocenica</i>	—	+	—	—	—
<i>M. japonica</i>	—	+	—	—	—
<i>M. varians</i> Agardh	—	+	—	—	—
<i>Alveolophora kbursevichiae</i>	+	—	—	—	—
<i>A. tscheremissinovae</i> (= <i>Miosira tscheremissinovae</i>)	+	—	—	—	+
<i>A. jouseana</i>	—	—	—	+	+
<i>A. areolata</i>	—	—	—	+	+
<i>Miosira antiqua</i>	—	—	—	—	+
Taxa counts by the source	10 taxa of 4 genera	12 taxa of one genus	8 taxa of one genus	14 taxa of 4 genera	15 taxa of 4 genera

Note: (+) taxon is present, (–) taxon is absent.

* Today, some species of the genus *Melosira* were transferred to *Aulacoseira* (Simonsen 1979).

The spellings of taxa are according to the authors.

crumbles constantly and reveals more and more layers of rocks. Consequently, within this period of time, the diatomologists could study the samples from different layers, which could be accumulated in different parts of the ancient lake (sublittoral, pelagic zone), and, possibly, even in different lakes. Pavlyutkin (2005), on the basis of the lithological-facies analysis, notes that, in the Middle Miocene time on this territory, there were lakes located at the floodplain of a large river (floodplain lakes) or at its delta (delta lakes). Nikolskaya (1952), Ivashinnikov (1978) argue that the Middle Miocene is a time of lacustrine transgression in this area. Other researchers believe that a basin similar to the present-day Khanka Lake was absent in the Miocene, and “lake baths” were in its place (Khudyakov et al. 1972, Pavlyutkin 1986, 2005). This is indicated by groups of facies of

floodplains, lakes and flowing lakes (Ablaev et al. 1994). Data reported by Jousé (1952) can be a prime example of the difference in diatom flora sampled from different habitats. Jousé has studied two sections located on the western shore of the lake. The sediments of one of the sections contained the *Melosira* representatives with thin valves and plenty of spines located along the margin of the valve; the rough silicified frustules of this genus were found in sediments of another section. Jousé has noted that the structure of forms with thin valves was very subtle, sometimes barely visible and close to the structure of modern *Melosira* (= *Aulacoseira*) *italica* Kützing including the following taxa: *Melosira italica* f. *miocenica* Jousé, *M. italica* var. *tenuissima* f. *miocenica* Jousé, *M. japonica* Pantocsek. Unfortunately, the thin-walled taxa discovered by Jousé cannot be compared either with fossil

or with modern ones, since there are only brief diagnoses without micrographs in the paper (Jousé 1952).

Strange as it may sound, the next problem complicating the comparative analysis of the Khanka diatom flora is the development of the instrumental base. The introduction of electron microscopy into the morphological studies after the 1970s made it possible to carry out a detailed study of the diatom valves ultrastructure and revolutionized the taxonomy of diatoms (Loseva 2002). As a result, a majority of the *Melosira* taxa specified in the Jousé and Moiseeva papers were transferred to another genus – *Aulacoseira*. *Melosira prae-islandica* (O. Müller) Jousé and *M. prae-distans* Jousé described by Jousé were included as synonymics of *Aulacoseira praegr anulata* var. *praeislandica* f. *praeislandica* (Simonsen) Moiseeva (Glezer et al. 1992).

So, *Melosira scabrosa* Østrup found by Moiseeva (1971) in the Miocene deposits of the Khanka Lake area was transformed in 1986 to *Paralia scabrosa* (Østrup) Moiseeva (Moiseeva 1986); currently, this taxon is *Ellerbekia arenaria* var. *teres* (Brim) Crawford (Glezer et al. 1992). According to the diagnosis and LM microphotographs, *Melosira scabrosa* (Moiseeva 1971) is characterized by short ridges up to 1/3 or 1/2 of the valve radius (7–9 in 10 µm), which is similar to the *Ellerbekia kochii* found by us. *E. kochii* differs from *E. arenaria* var. *teres* by a length of ridges on the valve face and the features of tube processes on the internal surface. The ridges on valve face of *E. kochii* are shorter than 1/3 of the valve radius, the number of ridges is 6–10 in 10 µm, original tube processes (2–3 in 10 µm) are formed by 4–6 ribs curved in spiral order, whereas valves of *E. arenaria* var. *teres* are characterized by short ridges up to 1/3 of the valve radius, 4–6 in 10 µm, the polyhedrous tube processes are formed by four, rarely, two ribs (Glezer et al. 1992). The number of ribs forming the polyhedrous tube processes can be counted only using SEM. Often, *E. kochii* has considered in *E. arenaria* var. *teres* volume (Glezer et al. 1992). Additional studies of *Ellerbekia* morphology are necessary for a correct diagnostics of taxa from this genus.

It is rather difficult to compare data on the quantitative participation of taxa in the diatom taphocenoses. It is associated with different systems of diatom counting used by the researchers. Jousé (1952) and Moiseeva (1971) have used the Visloukh grading scale with following categories: separately, rare, not infrequently, frequently, over and over again and in the mass. At present, the diatomologists calculate the percentage of diatom taxa in the taphocenoses. This makes possible to understand the ratio of taxa in the fossil community. Thus, even a short list of problems shows that further study of the Khanka diatom flora is needed.

Nevertheless, owing to the accumulated data and detailed examination carried out by us, the Middle Miocene diatom centric flora was sufficiently characterized in order to assess its image and perform comparison with the even-aged diatom floras studied from continental deposits of other regions. Unfortunately, there is no one continuous sequence of Neogene (23–1.8 Ma) continental deposits of the Far East of Russia. There are only separated sections located at different points of the region which makes their correlation difficult. In addition, the continental diatoms,

in comparison with the marine ones, have high rates of speciation due to the diversity of terrestrial biotopes and variability of ecological parameters. They are also sensitive to fluctuations of the environmental conditions which results in a wide range of phenotypic variability. The listed diatom features make them a good tool for reconstructing the paleoenvironmental parameters. However, they complicate the correlation of faciesly heterogeneous strata, since deposits may contain the diatom complexes with incommensurable ecological structures (Likhacheva et al. 2009). When identifying the diatom complexes and their using for local and regional correlation, it is necessary to be founded, first of all, on the diatom evolution. Undoubtedly, this process should be considered in combination with the geological and paleogeographical events fixed on this territory. The understanding of these regularities increases validity of constructions based on analysis of floras. The established phasing provides the possibility to creating the zonal stratigraphic continental scales. The maximal certainty is characteristic of the biostratigraphic zonal scales where the changes of the paleoecosystems with close living environment and, in the aggregate, constituting the chronology of the paleobiosphere variations during specified time interval are reflected. At present, the zonal biostratigraphic scales for the Neogene of Primorye (Likhacheva et al. 2009) and Russian Far East (Moiseeva 1995) were successfully developed; the age distribution of diatom complexes was determined. This allows us to compare the Middle Miocene centric flora of the Khanka Depression with the even-aged complexes from other regions of Far East.

The Middle Miocene centric diatom flora of the Khanka Depression has a particular similarity to the even-aged floras from other regions of Primorye Territory: the Tony Lake area (basin of the Samarga River) and the basin of Suputinka River (Moiseeva 1971), and also from Baikal region (Popova et al. 1989, Chernyaeva et al. 2007, Enikeev 2008), and eastern China (Li et al. 2010). In floras from all the areas, the centric taxa dominate. *Melosira praeislandica* f. *praeislandica* (= *Aulacoseira praegr anulata* var. *praeislandica* f. *praeislandica*) dominates in deposits of Tony Lake area and the basin of Suputinka River. *Melosira scabrosa* (= *Ellerbekia kochii*) dominates in the diatom complex from the basin Supputinka River. *Melosira undulata* is absent in the deposits from the Tony Lake area. This is the difference between these floras and the Khanka flora. The *Melosira* (= *Aulacoseira*) genus has dominated in the Middle Miocene deposits of the Chara Basin located in the eastern part of the Baikal-Stanovoy Upland (Enikeev 2008). In these deposits, there are following taxa from the genus *Melosira*: *Melosira praedistans* Jousé, *Melosira* sp. (aff. *canadensis*), *M. distans* (Ehrenberg) Kützing, *M. distans* var. *alpigena* Grunow, *M. italica* (Ehrenberg) Kützing, *M. aff. praegr anulata* Jousé. *Paralia scabrosa* (Østrup) Moiseeva is subdominant. The diatom photos are absent in the Enikeev article, therefore, we can only assume that, judging by mention of name *Melosira* genus renamed at present as *Aulacoseira* and those renamings which we fore mentioned, in the Middle Miocene complex of the Chara Basin, the *Aulacoseira praegr anulata* var. *praeislandica* f. *praeislandica* and *Ellerbekia*

kochii were found. *Aulacoseira praegr anulata* var. *praegr anulata*, *A. praegr anulata* var. *praeislandica*, *Pseudoaulacosira moiseeviae* (Lupikina) Lupikina & Khursevich ex Usoltseva & Houk dominate in Dzhilinskaya Depression deposits of Western Transbaikalia; *Alveolophora jouseana* (Moiseeva) Moiseeva, *Melosira undulata*, *Aulacoseira baicalensis* (K. Meyer) Simonsen take lesser part (Chernyaeva et al. 2007). *Aulacoseira* cf. *distans* (Ehrenberg) Simonsen and *Melosira youngi* Skvortsov were found in the Middle Miocene Shanwang Formation, Shandong Province, East China (Li et al. 2010). Thus, this may be regarded as a wide distribution of centric taxa in the Middle Miocene water bodies of the entire Far East.

It is interesting to note that the freshwater centric diatoms were quite widely distributed in the Miocene of Europe (Khursevich 2007), Turkey (Servant-Vildary et al. 1988), USA, Oregon (Ledgerwood & Tassell 2006) and so on. These are also the representatives of the genus *Aulacoseira*. The record of the genus *Aulacoseira* goes back not less than 45 million years (Khursevich 1995, Krebs 1994), and habitation of this genus in the Tertiary lacustrine deposits is natural occurrence (Haworth & Sabater 1993, Khursevich 1995). Probably, such wide occurrence and dominance of the representatives of this taxon in the Middle Miocene is related to the particular evolutionary stage of development of diatoms when the maximum generic and species diversity of the freshwater centric diatoms is observed. Likely, the quick evolution of this genus is a result of the global climatic phenomena, particularly, climatic optimum of the Middle Miocene which followed on the cold spell of Oligocene. According to the data of paleobotany (Bondarenko et al. 2019), the lukewarm conditions were established for the Late Oligocene (24.0 ± 3.0 mil. years) in Primorye. According to the data for diatom algae (Likhacheva et al. 2009), the warming has begun in the Miocene and reached a peak in its middle. The Middle Miocene was warmest within a period of 30 million years (Zachos et al. 2001).

CONCLUSION

The freshwater centric diatoms of the *Aulacoseira*, *Alveolophora*, *Ellerbeckia* and *Melosira* genera from the Middle Miocene deposits of Primorye (Far East of Russia) were studied in detail using the light and scanning electron microscopes.

As a result, the considerable morphological variability of predominant representatives of the genus *Aulacoseira* was established. The measurements of the valve heights and diameters of *Aulacoseira* taxa have shown that *A. praegr anulata* var. *praeislandica* f. *praeislandica* are characterized by high variability of these parameters; *A. praegr anulata* var. *praegr anulata* f. *praegr anulata* has the constant characteristics; *A. praegr anulata* var. *praeangustissima* f. *praeangustissima* has the constant heights of valves.

The diagnoses of *Alveolophora tscheremissinovae*, *Ellerbeckia kochii* and *Melosira undulata* were supplemented.

Unfortunately, we don't find the centric taxa previously identified by others researchers such as *Alveolophora jouseana* (Moiseeva) Moiseeva, *Alveolophora areolata* Moiseeva (Pavlyutkin et al. 2004, Likhacheva et al. 2009), oval *Miosira antiqua* (Moiseeva) Khursevich and taxa of *Actinocyclus* Ehrenberg, *Mesodictyon* E.C. Theriot & J.P. Bradbury genera (Likhacheva

et al. 2009). Additional researches are necessary for complete evaluation of the taxonomic composition. New data from Middle Miocene-Holocene diatomaceous sediments of the Khanka Depression need to be investigated. The samples selected from the continuous sections that preserve the consistent lake history will provide the additional data for objective paleogeographic reconstruction and the diatom flora evolution in the past.

ACKNOWLEDGEMENTS

This study were supported by the Programs of Fundamental Researches of the Federal scientific center of the East Asia terrestrial biodiversity FEB RAS (№ 0267-2019-0015), of the Limnological Institute SB RAS (№ 0279-2021-0008) and RFBR, project № 19-04-00746. The authors appreciate of Vitaly Kazarin for helping with the scanning electron microscope.

LITERATURE CITED

- Ablaev, A.G., S.M. Tashchi & I.V. Vassiliev 1994. *Miocene of the Khanka Depression of the West Primorye*. Dalnauka, Vladivostok, 168 pp. (in Russian). [Аблаев А.Г., Ташчи С.М., Васильев И.В. 1994. Миоцен Ханкайской впадины Западного Приморья. Владивосток: Дальнаука. 168 с.].
- Bondarenko, O.V., N.I. Blokhina & T. Utescher 2019. Major plant biome changes in the Primorye Region (Far East of Russia) during the Paleogene. *Botanica Pacifica* 8(1):3–15.
- Chernyaeva, G.P., N.A. Lyamina, S.V. Rasskazov, I.N. Rezanov & V.V. Savinova 2007. Biostratigraphy and deposition environments of the Middle-Late Miocene volcanosedimentary section in the Dzhilinda basin, Western Transbaikalia. *Russian Geology and Geophysics* 48(4):361–370.
- Chernyaeva, G.P. & A.I. Moiseeva 2003. Diatom assemblages from Upper Cenozoic sediments of the Chara depression (Transbaikalia Region). *Doklady Earth Sciences* 391(5):659–662.
- Chernyaeva, G.P. & S.M. Popova 1993. The discovery of diatoms in the lower Tankhoi suite, South Baikal basin. *Geografiya i Prirodnye resursy* 3:168–171 (in Russian). [Черняева Г.П., Попова С.М. 1993. О первой находке диатомовых водорослей в низах танхойской свиты (Южно-Байкальская впадина) // География и природные ресурсы. № 3. С. 168–171].
- Cherepanova, M.V., J.A. Snyder & J. Brigham-Grette 2007. Diatom stratigraphy of the last 250ka at El'gygytyn Lake, northeast Siberia. *Journal of Paleolimnology* 37:155–162.
- Cherepanova, M.V., M.V. Usoltseva, V.S. Pushkar & Yu.F. Dubrovina 2010. Morphogenesis in *Cyclotella ocellata* complex from Lake El'gygytyn (Chukchi Peninsula) during the Pleistocene–Holocene. *Paleontological Journal* 44(10):1252–1261.
- Cohen, K.M., D.A.T Harper & P.L. Gibbard. 2021. *ICS International Chronostratigraphic Chart 2021/01*. International Commission on Stratigraphy, IUGS. Available from: <https://www.stratigraphy.org/>. Last accessed 18.06.2021.
- Douglas, M.S.V. & J.P. Smol 1999. Freshwater diatoms as indicators of environmental change in the High Arctic. In: *The Diatoms: Applications for the Environmental and Earth Sciences, 2nd edition* (J.P. Smol, ed.), pp. 227–244, Cambridge University Press, Cambridge.
- Dubrovina, Yu.F., M.V. Usoltseva, V.S. Pushkar & O.Yu. Likhacheva 2011. The morphology of dominant species *Tetracyclus* Ralfs from Miocene sediments of the Novokachalinskaya Suite (Primorye, Russia). In: *Diatoms: morpho-*

- logy, systematics, floristics, ecology, paleogeography, biostratigraphy: proceedings of the XII International scientific conference of diatomologists in tribute of the 120-years anniversary of A.I. Proshkina-Lavrenko (Moscow, September 19-24, 2011) (V.M. Gavrilov, ed.), pp. 233–235, University books, Moscow (in Russian). [Дубровина Ю.Ф., Усольцева М.В., Пушкар В.С., Лихачева О.Ю. 2011. Морфология доминирующих видов *Tetracyclus* Ralfs из миоценовых отложений новокачалинской свиты (Приморье, Россия) // Диатомовые водоросли: морфология, систематика, флористика, экология, палеогеография, биостратиграфия: материалы XII Международной научной конференции диатомологов посвященной 120-летию со дня рождения А.И. Прошкиной-Лавренко (19–24 сентября 2011 г., Москва) / ред. Гаврилов В.М. Москва: Университетская книга. С. 233–235].
- Dubrovina, Yu., M. Usoltseva, V. Pushkar & O. Likhacheva 2014. *Tetracyclus chankaicus*, a new species from Miocene sediments of the Khanka Lake, Primorye, Russia. *Diatom Research* 29(2):111–118.
- Ehrenberg, G.C. 1840. Charakteristik von 274 neuen Arten von Infusorien. In: *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich-Preussischen Akademie der Wissenschaften zu Berlin*, s. 197–219, Berlin.
- Enikeev, F.I. 2008. The Late Cenozoic of Northern Transbaikaliya and paleoclimate of Southern East Siberia. *Russian Geology and Geophysics* 49(8):602–610.
- Ershova, T.B. 2015. *Statistics, part 2*. Izdatel'stvo AmGPU, Komsomolsk-na-Amure, 74 pp. (in Russian). [Ершова Т.Б. 2015. Статистика. В 2 частях. Ч. 2. Комсомольск-на-Амуре: Изд-во АМГПУ. 74 с.].
- Flower, B.P. & J.P. Kennett 1994. The Middle Miocene climatic transition: East Antarctic ice sheet development, deep ocean circulation and global carbon cycling. *Palaeogeography, Palaeoclimatology, Palaeoecology* 108(3–4): 537–555.
- Fritz, S.C., S. Juggins & R.W. Battarbee 1993. Diatom assemblages and ionic characterization of lakes of the northern great-plains, North-America: a tool for reconstructing past salinity and climate fluctuations. *Canadian Journal of Fisheries and Aquatic Sciences* 50:1844–1856.
- Fritz, S.C., S. Juggins, R.W. Battarbee & D.R. Engstrom 1991. Reconstruction of past changes in salinity and climate using a diatom-based transfer function. *Nature* 352:706–708.
- Glezer, Z.I., I.V. Makarova, A.I. Moiseeva & A.V. Nikolaev 1992. *The diatoms of the USSR. Fossil and recent, vol. 2, issue 2. Stephanodiscaceae, Ectodictyonaceae, Paraliaceae, Radialiplicataceae, Pseudopodosiraceae, Trochosiraceae, Melosiraceae, Aulacosiraceae*, Nauka, Saint-Petersburg. 125 pp. (in Russian). [Глезер З.И., Макарова И.В., Моисеева А.И., Николаев А.В. 1992. Диатомовые водоросли СССР (ископаемые и современные). Т. II, вып. 2. Stephanodiscaceae, Ectodictyonaceae, Paraliaceae, Radialiplicataceae, Pseudopodosiraceae, Trochosiraceae, Melosiraceae, Aulacosiraceae. Санкт-Петербург: Наука. 125 с.].
- Guiry, M.D. & G.M. Guiry 2021. *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Available from: <http://www.algaebase.org/>. Last accessed 18.06.2021.
- Hassan, A., M. Usoltseva, S. Rasskazov, I. Chuvashova & L. Titova 2019. First study of fossil diatom flora from Middle Miocene-Lower Pliocene lacustrine sediments in Barguzin Valley, Baikal Rift Zone. *Quaternary International* 524:24–30.
- Haworth, E.Y. & S. Sabater 1993. A new Miocene *Aulacoseira* species in diatomite from the ancient lake in La Cerdanya (NE Spain). *Nova Hedwigia* 106:227–242.
- Hayashi, T., M. Saito-Kato & Y. Tanimura 2012. *Actinocyclus nipponicus* sp. nov. and *A. bradburyi* sp. nov. (Bacillariophyta) from Miocene lacustrine sediments of the proto-Japan Sea. *Phycologia* 51(1):98–112.
- Haywood, A.M., J.L. Smellie, A.C. Ashworth, D.J. Cantrill, F. Florindo, M.J. Hambrey, D. Hill, C.-D. Hillenbrand, S.J. Hunter, R.D. Larter, C.H. Lear, S. Passchier & R. van de Wal 2009. Middle Miocene to Pliocene history of Antarctica and the Southern Ocean. In: *Developments in Earth & Environmental Sciences «Antarctic Climate Evolution», vol. 8* (F. Florindo & M. Siegert, eds.), pp. 403–461, Elsevier Science & Technology, England.
- Hëribaud, J. 1893. *Les Diatomées fossiles d'Auvergne*. Clemont-Ferrand, Paris, 255 pp.
- Hustedt, F. 1930. Die Kieselalgen Deutschlands, Österreichs und der Schweiz: mit Berücksichtigung der übrigen Länder Europas sowie der angrenzenden Meeresgebiete. Teil. 1. In: *Dr. L. Ra-benhorst's Kryptogamenflora von Deutschland, der Schweiz, Bd 7, Reprint 1977* (O. Koeltz, ed.), ss. 557–816, Koenigstein, West Germany.
- Ivashinnikov, Yu.K. 1978. *Paleogeomorphology of the depression morphostructures in the south of the Far East*. Nauka, Moscow, 131 pp. (in Russian). [Ивашинников Ю.К. 1978. Палеогеоморфология депрессионных морфоструктур юга Дальнего Востока. Москва: Наука. 131 с.].
- Jiang, H. & Z. Ding 2008. A 20 Ma pollen record of East-Asian summer monsoon evolution from Guyuan, Ningxia, China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 265:30–38.
- Jousé, A.P. 1952. On the history of diatom flora of the Khanka Lake. *Trudy Instituta geografii AN SSSR «Materialy po geomorfologii i paleogeografii SSSR»* 6(51):226–252 (in Russian). [Жузе А.П. 1952. К истории диатомовой флоры озера Ханка // Труды института географии АН СССР «Материалы по геоморфологии и палеогеографии СССР». Т. 6, Вып. 51. С. 226–252].
- Kennett, J. 1982. *Marine geology*. Prentice-Hall, New Jersey, 813 pp.
- Khudyakov, I.G., E.P. Denisov, A.M. Korotkiy, A.A. Kulakov, R.F. Nikonova & E.I. Chernobrovina 1972. *The history of the development of the relief of Siberia and the Far East. South of the Far East*. Nauka, Moscow, 423 pp. (in Russian). [Худяков И.Г., Денисов Е.П., Короткий А.М., Кулаков А.А., Никонова Р.Ф., Чернобровина Е.И. 1972. История развития рельефа Сибири и Дальнего Востока. Юг Дальнего Востока. Москва: Наука. 423 с.].
- Khursevich, G.K. 1994. Morphology and taxonomy of some centric diatom species from the Miocene sediments of the Dzhilinda and Tunkin hollows. In: *Memoirs of the California Academy of Sciences, vol. 17: proceedings of the 11th International Diatom Symposium (San Francisco, California, USA, August 12–17, 1990)* (J.P. Kocielek, ed.), pp. 269–280, California Academy of Sciences, San Francisco.
- Khursevich, G.K. 1995. Evolution of freshwater centric diatoms within the Eurasian continent. In: *Proceedings of the Thirteenth International Diatom Symposium (Maratea, Italy, September 1–7, 1994)* (D. Marino & M. Montresor, eds), pp. 507–520, Biopress Limited, Bristol.
- Khursevich, G.K. 2007. Evolution stages of freshwater centric diatoms in the Cenozoic within the Euroasian continent. In: *Morphology, cell biology, ecology, floristry and history of the development of diatoms: proceedings of the X International Conference of Diatomologists of the CIS countries (Minsk, Belarus, September 09–14, 2007)* (G.K. Khursevich, ed.) pp. 225–228, BGPU, Minsk (in Russian with English summary). [Хурсевич Г.К. 2007. Этапность в эволюции прес-

- новодных центрических диатомей на протяжении кайнозоя на Евразийском континенте // Морфология, клеточная биология, экология, флористика и история развития диатомовых водорослей: материалы X Международной конференции диатомологов стран СНГ (09–14 сентября 2007 г., Минск, Беларусь) / ред. Г.К. Хурсевич. Минск: БГПУ. С. 225–228].
- Khursevich, G.K. & J.P. Kociolek 2012. A preliminary, worldwide inventory of the extinct, freshwater fossil diatoms from the orders Thalassiosirales, Stephanodiscales, Paraliales, Aulacoseirales, Melosirales, Coscinodiscales and Biddulphiales. *Nova Hedwigia* 141:315–364.
- Khursevich, G.K., A.A. Prokopenko, S.A. Fedenya, L.I. Tkachenko & D.F. Williams 2005. Diatom biostratigraphy of Lake Baikal during the past 1.25 Ma: new results from BDP-96-2 and BDP-99 drill cores. *Quaternary International* 136(1):95–104.
- Krammer, K. & H. Lange-Bertalot 1991. Bacillariophyceae 3. Teil: Centrales, Fragilariaceae, Eunotiaceae. In: *Süßwasserflora von Mitteleuropa, b. 2(3)* (H. Ettl, J. Gerloff, H. Heynig, D. Mollenhauer, eds.), 33. 1–576, Gustav Fisher Verlag, Stuttgart, Germany.
- Krapp, M. & J.H. Jungclaus 2011. The Middle Miocene climate as modelled in an atmosphere-ocean-biosphere model. *Climate of the Past* 7:1169–1188.
- Krebs, W.N. 1994. The biochronology of freshwater planktonic diatom communities in western North America. In: *Memoirs of the California Academy of Sciences, vol. 17: proceedings of the 11th International Diatom Symposium (San Francisco, California, USA, August 12–17, 1990)* (J.P. Kociolek, ed.), pp. 485–499, California Academy of Sciences, San Francisco.
- Kürschner, W.M., Z. Kvaček & D.L. Dilcher 2008. The impact of Miocene atmospheric carbon dioxide fluctuations on climate and the evolution of terrestrial ecosystems. *PNAS* 105(2):449–453.
- Kützing, F.T. 1844. *Die Kieselalgen. Bacillarien oder Diatomeen*. Nordhausen. 152 ss.
- Kuzmin, M.I., G.K. Khursevich, A.A. Prokopenko, S.A. Fedenya & E.B. Kabanov 2009. *Centric diatoms in Lake Baikal during the Late Cenozoic: morphology, systematics, stratigraphy, and stage of development (Based on the Deep Cores of the Baikal Drilling Project)*. GEO, Novosibirsk, 374 pp. (in Russian). [Кузьмин М.И., Хурсевич Г.К., Прокопенко А.А., Феденя С.А., Кабанов Е.Б. 2009. Центрические диатомовые водоросли позднего кайнозоя озера Байкал. Новосибирск: GEO, 2009. 374 с.].
- Ledgerwood, R. & J. Van Tassell 2006. Stratigraphy and age the late Miocene sediments and volcanic deposits along the Baker-Copperfield Highway between Banta road and the Love Ranch, Keating, Oregon. *Eastern Oregon Geology* 3:52–70.
- Lewis, A.R., D.R. Marchant, A.C. Ashworth, L. Hedenäs, S.R. Hemming, Jesse V. Johnson, M.J. Leng, M.L. Machlus, A.E. Newton, J.I. Raine, J.K. Willenbring, M. Williams & A.P. Wolfe 2008. Mid-Miocene cooling and the extinction of tundra in continental Antarctica. *PNAS* 105(31):10676–10680.
- Li, Y.-M., D.K. Ferguson, Y.-F. Wang & C.-S. Li 2010. Paleoenvironmental inferences from diatom assemblages of the middle Miocene Shanwang Formation, Shandong, China. *Journal of Paleolimnology* 43:799–814.
- Likhacheva, O.Yu. 2013. *Main geological events of the Neogene Southern Primorye (diatom analysis)*. PhD dissertation theses. Far East Geological Institute FEB RAS, Vladivostok, 26 pp. (in Russian). [Лихачева О.Ю. 2013. Основные геологические события неогена юга Приморья (диатомовый анализ): автореферат дис. ... канд. геол.-мин. наук. Владивосток: Дальневосточный Геологический институт ДВО РАН. 26 с.].
- Likhacheva, O.Yu., V.S. Pushkar, M.V. Cherepanova & B.I. Pavlyutkin 2009. Zonal diatom scale and the major geobiological events of Primorye Neogene. *Vestnik DVO RAN* 4:64–72 (in Russian with English summary). [Лихачева О.Ю., Пушкар В.С., Черепанова М.В., Павлюткин Б.И. 2009. Зональная диатомовая шкала и основные геобиологические события неогена Приморья // Вестник ДВО РАН. № 4. С. 64–72].
- Likhoshway, E.V. & R.M. Crawford 2001. The rimoportula - a neglected feature in the systematics of Aulacoseira. In: *Abstracts of the 16th International Diatom Symposium (Athens, Greece, August 28–September 1, 2000)* (A. Economou-Amilli, ed), p. 64, Amvrosiou Press, Athens.
- Likhoshway, E.V., G.V. Pomazkina & T.A. Nikiteeva 1997. Centric diatoms of the Middle Miocene in the Baikal rift zone (Tunka basin). *Geologiya i Geofizika* 38(9):1445–1452 (in Russian with English summary). [Лихошвай Е.В., Помазкина Г.В., Никитеева Т.А. 1997. Центрические диатомовые водоросли из миоценовых отложений Байкальской рифтовой зоны (Тункинская котловина) // Геология и геофизика. Т. 38, № 9. С. 1445–1452].
- Loseva, E.I. 1981. Some common fossil diatoms from the east European plain (the Kama River Basin) and the Massif Central. In: *Taxonomy, Morphology, Ecology, Biology: proceedings of the Sixth Symposium on Recent and Fossil Diatoms (Budapest, Hungary, September 1–5, 1980)* (R. Ross, ed), pp. 75–80, Koeltz Publishing, Koenigstein.
- Loseva, E.I. 2002. *Beautiful invisibles*. Ural Branch of RAS, Ekaterinburg, 146 pp. (in Russian with English summary). [Лосева Э.И. 2002. Прекрасные невидимки. Екатеринбург: УрО РАН. 146 с.].
- Lupikina, E.G. 1991. New family Ellerbeckiaceae Lupik. (Bacillariophyta). *Algologia* 1(2):58–64.
- Manoylov, K.M., N. Ognjanova-Rumenova & R.J. Stevenson 2009. Morphotype variations in subfossil diatom species of Aulacoseira in 24 Michigan Lakes, USA. *Acta Botanica Croatica* 68(2):401–419.
- Mats, V.D. 2013. Late Cretaceous and Cenozoic stratigraphy of the Baikal Rift sediments. *Stratigraphy and Geological Correlation* 21(6):637–651.
- Moiseeva, A.I. 1971. Atlas of Neogene Diatom Algae of Primorsky region. In: *Trudy VSEGEI. Novaya seriya, vol. 171* (Z.I. Glezer, ed.), 152 pp., Nedra, Leningrad (in Russian). [Моисеева А.И. 1971. Атлас неогеновых диатомовых водорослей Приморского края // Труды ВСЕГЕИ. Новая серия / под ред. З.И. Глезер. Ленинград: Недра, Т. 171. 152 с.].
- Moiseeva, A.I. 1986. Revision of the systematic position of freshwater species of the genus *Melosira* Ag. s. l. (Bacillariophyta). In: *Sbornik Nauchnykh Trudov "Aktual'nye voprosy sovremennoi paleoalgologii"* (B.S. Sokolov, ed.), pp. 70–73, Naukova dumka, Kiev (in Russian). [Моисеева А.И. Ревизия систематического положения пресноводных видов рода *Melosira* Ag. s. l. (Bacillariophyta) // Сборник научных трудов «Актуальные вопросы современной палеоальгологии» / Под ред. Б.С. Соколова. Киев: Наукова думка. С. 70–73].
- Moiseeva, A.I. 1995. Diatom zonation of Neogene continental deposits of the Russian Far East. *Stratigrafiya. Geologicheskaya Korrelyatsiya* 3(5):92–103 (in Russian). [Моисеева А.И. 1995. Расчленение континентальных отложений неогена Дальнего Востока по диатомеям //

- Стратиграфия. Геологическая корреляция. Т.3, № 5. С. 92–103].
- Moiseeva, A.I. & S.I. Genkal 1987. On fresh water species of the genus *Paralia* (Bacillariophyta). *Botanicheskii Zhurnal* 72 (11):1500–1504 (in Russian). [Моисеева А.И., Генкал С.И. 1987. О пресноводных видах рода *Paralia* (Bacillariophyta) // Ботанический журнал. Т. 72, № 11. С. 1500–1504].
- Moiseeva, A.I. & E.I. Tsarko 1990. Diatoms flora stages of development and detailed stratigraphy of the Neogene in south of the Russian Far East. In: *New data on stratigraphy of the Far East and the Pacific Ocean* (A.G. Ablaev, ed.), pp. 68–78, Far East Branch of the Russian Academy of Sciences, Vladivostok (in Russian). [Моисеева А.И., Царько Е.И. 1990. Этапы развития флоры диатомовых водорослей и детальная стратиграфия неогена юга Дальнего Востока // Новые данные по стратиграфии Дальнего Востока и Тихого океана / под ред. А.Г. Аблаева. Владивосток: ДВО РАН СССР. Р. 68–78].
- Nikolaev, V.A. & D.M. Harwood 2001. Diversity and classification of centric diatoms. In: *Proceeding of the 16th International Diatom Symposium (Athens, 25 August – 1 September 2000)* (A. Economou-Amilli, ed.), pp. 127–152, University of Athens, Athens.
- Nikolskaya, V.V. 1952. Some data on the paleogeography of Lake Khanka. *Trudy instituta geografii AN SSSR "Materialy po geomorfologii i paleogeografii SSSR"* 6(51):215–226 (in Russian). [Никольская В.В. 1952. Некоторые данные по палеогеографии озера Ханка // Труды института географии АН СССР «Материалы по геоморфологии и палеогеографии СССР». Т. 6, Вып. 51. С. 215–226].
- Pantocsek, J. 1905. *Beiträge zur Kenntniss der fossilen Bacillarien Ungarns. B. 3: Süßwasser Bacillarien*. Buchdruckerei von Julius Platzko, Nagytapolcsány Topo, 118 ss.
- Pavlyutkin, B.I. 1986. *Geological development of the Razdolnaya-Khanka Depression in the Late Cenozoic*. PhD dissertation theses. Institute of Geology and Geophysics SB AN USSR, Novosibirsk, 16 pp. (in Russian). [Павлюткин Б.И. 1986. Геологическое развитие Раздольненско-Ханкайской впадины в поздне кайнозое: автореферат. дис. ... канд. геол.-мин. наук. Новосибирск: Институт геологии и геофизики СО АН СССР. 16 с.].
- Pavlyutkin, B.I. 2005. *The Mid-Miocene Khanka flora of the Primorye*. Dalnauka, Vladivostok, 216 pp. (in Russian with English summary). [Павлюткин Б.И. 2005. Среднемиоценовая ханкайская флора Приморья. Владивосток: Дальнаука. 216 с.].
- Pavlyutkin, B.I., S.S. Ganzey, V.S. Pushkar & T.I. Petrenko 1993. Paleobotany characteristics and radiometric dating of the Neogene deposits of southern Primorye. *Stratigrafiya. Geologicheskaya Korrelyatsiya* 1(6):40–47 (in Russian with English summary). [Павлюткин Б.И., Ганзей С.С., Пушкар В.С., Петренко Т.И. 1993. Палеоботаническая характеристика и радиометрическое датирование неогеновых отложений Южного Приморья // Стратиграфия. Геологическая корреляция. Т. 1, № 6. С. 40–47].
- Pavlyutkin, B.I., V.S. Pushkar, M.V. Cherepanova & T.I. Petrenko 2004. The Miocene stratigraphy problems of the Khanka depression of Primorye, Russian Far East. *Tikhookeanskaya Geologiya* 23(4):73–85. [Павлюткин Б.И., Пушкар В.С., Черепанова М.В., Петренко Т.И. 2004. Проблемы стратиграфии миоцена Приханкайской впадины (Дальний Восток России) // Тихоокеанская геология. Т. 23, № 4. С. 73–85].
- Polyakova, Ye.I. & I.D. Danilov 1989. The Miocene of the Far North of Western Siberia. *Doklady AN SSSR* 308(2):428–431 (in Russian). [Полякова Е.И., Данилов И.Д. 1989. Миоцен крайнего севера Западной Сибири (полуостров Ямал) // Доклады АН СССР. Т. 308, № 2. С. 428–431].
- Popova, C.M., V.D. Mats, G.P. Chernyaeva, M.K. Shimaraeva, A.A. Kulchitskiy, G.A. Vorobieva, V.M. Klimanova, E.E. Kononov, A.Ya. Kravchinsky, N.V. Kulagina, F.I. Lazo, L.A. Orlova, V.A. Panychev, A.M. Sizikov & I.V. Shibanova 1989. *Paleolimnological reconstructions: Baikal Rift zone*. Nauka, Novosibirsk, 109 pp. (in Russian with English summary). [Попова С.М., Мац В.Д., Черняева Г.П., Шимараева М.К., Кульчицкий А.А., Воробьева Г.А., Климанова В.М., Кононов Е.Е., Кравчинский А.Я., Кулагина Н.В., Лазо Ф.И., Орлова Л.А., Паньчев В.А., Сизиков А.М., Шибанова И.В. 1989. Палеолимнологические реконструкции: Байкальская рифтовая зона. Новосибирск: Наука. 111 с.].
- Proshkina-Lavrenko, A.I. (chief ed.) 1974. *The diatoms of the USSR. Fossil and recent*. Nauka, Leningrad, 403 pp. (in Russian). [Диатомовые водоросли СССР (ископаемые и современные) / отв. ред. А.И. Прошкина-Лавренко. 1974. Ленинград: Наука. 403 с.].
- Pushkar, V.S. & O.Yu. Likhacheva 2007. Khanka diatom flora of Primorye and its position in the regional stratigraphic scheme of the Neogene. In: *Morphology, cell biology, ecology, floristry and history of the development of diatoms: proceedings of the X International Conference of Diatomologists of the CIS countries (Minsk, Belarus, September 09–14, 2007)* (G.K. Khursevich, ed.), pp. 215–218, BGPU, Minsk (in Russian with English summary). [Пушкар В.С., Лихачева О.Ю. 2007. Ханкайская диатомовая флора Приморья и ее положение в региональной стратиграфической схеме неогена // Морфология, клеточная биология, экология, флористика и история развития диатомовых водорослей: материалы X Международной конференции диатомологов стран СНГ (09–14 сентября 2007 г., Минск, Беларусь) / ред. Г.К. Хурсевич. Минск: БГПУ. С. 215–218].
- Rasskazov, S.V., N.A. Lyamina, G.P. Chernyaeva, I.V. Lusina, A.F. Rudnev & I.N. Resanov 2007. *Cenozoic stratigraphy of the Vitim Plateau: phenomenon of long rifting in the south of East Siberia*. GEO, Novosibirsk, 193 pp. (in Russian). [Рассказов С.В., Лямина Н.А., Черняева Г.П., Лузина И.В., Руднев А.Ф., Резанов И.Н. Стратиграфия кайнозоя Витимского плоскогорья: феномен длительного рифтогенеза на юге Восточной Сибири. Новосибирск: ГЕО. 193 с.].
- Rasskazov, S.V., N.A. Lyamina, I.V. Luzina, G.P. Chernyaeva, I.S. Chuvashova & M.V. Usoltseva 2014. Sediments in the Tertiary Tankhoi field, South Baikal basin: stratigraphy, correlation and structural transformations in the Baikal regions. *Geodynamics & Tectonophysics* 5(4):993–1032.
- Reid, M. 2005. Diatom-based models for reconstructing past water quality and productivity in New Zealand lakes. *Journal of Paleolimnology* 33:13–38.
- Round, F.E., R.M. Crawford & D.G. Mann 1990. *The diatoms: biology and morphology of the genera*. Cambridge University Press, Cambridge, 747 pp.
- Royden, L.H., B.C. Burchfiel & R.D. van der Hilst 2008. The geological evolution of the Tibetan plateau. *Science* 321:1054–1058.
- Saito-Kato, M., T. Hayashi & T. Tanaka 2011. *Tetracyclus radialis* (Bacillariophyta), a new fossil species from Miocene freshwater sediments in the Japan Sea. *Phytotaxa* 24:51–57.
- Schoell, M., S. Schouten, J.S. Sinninghe Damsté, J.W. de Leeuw & R.E. Summons 1994. A molecular organic carbon isotope record of Miocene climate changes. *Science*

- 263(5150):1122–1125.
- Shevenell, A.E. & J.P. Kennett 2004. Paleoceanographic change during the Middle Miocene Climate Revolution: an Antarctic stable isotope perspective. In: *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change Between Australia and Antarctica Geophysical Monograph, Series 151* (N.F. Exon, J.P. Kennett & M.J. Malone, eds.), pp. 235–251, American Geophysical Union, Washington.
- Servant-Vildary, S., J.C. Paicheler & B. Selemin 1988. Miocene Lacustrine diatoms of Turkey. In: *Proceeding of the 9th Diatom Symposium (Bristol, United Kingdom, August 24–30, 1986)* (F.E. Round, ed.) pp.165–180, Biopress Ltd., Bristol.
- Simonsen, R. 1979. The Diatom system: ideas on phylogeny. *Bacillaria* 2:9–71.
- Stoermer, E.F. & J.P. Smol 1999. *The diatoms: applications for the environmental and earth sciences*. University Press, Cambridge, 686 pp.
- Strelnikova, N.I. & I.B. Tsoy (eds.) 2008. *The diatoms of Russia and adjacent countries. Fossil and recent, vol. 2, issue 5*. Saint Petersburg University press, Saint Petersburg, 171 pp. (in Russian with English summary). [Диаомовые водоросли России и сопредельных стран. Ископаемые и современные / под. ред. Н.И. Стрельниковой, И.Б. Цой. 2008. Санкт-Петербург: Издательство Санкт Петербургского университета. Т. 2, вып. 5. 171 с.]
- Tanai, T. & K. Huzioka 1967. Climatic implications of Tertiary floras in Japan. In: *Tertiary correlation and climatic changes in the Pacific* (K. Hatai, ed.), pp. 89–94, Sasaki, Sendai.
- Tanaka, H. & T. Nagumo 2011a. *Aulacoseira bouki*, a new Early Miocene freshwater diatom from Hiramaki Formation, Gifu Prefecture, Japan. *Diatom* 26(2):161–165.
- Tanaka, H. & T. Nagumo 2011b. *Aulacoseira iwakiensis* sp. nov., a new elliptical *Aulacoseira* species from an Early Miocene sediment, Japan. *Diatom* 27:1–8.
- Tanaka, H. & T. Nagumo 2014. *Actinocyclus hiramakiensis*, a new freshwater diatom from Early Miocene sediment, Gifu Prefecture, Japan. *Diatom* 30:157–167.
- Tanaka, H., T. Nagumo & F. Akiba 2008. *Aulacoseira hachiyaensis* sp. nov., a new Early Miocene freshwater fossil diatom from Hachiya Formation, Japan. In: *Proceedings of the Nineteenth International Diatom Symposium (Irkutsk, Russia, August 28–September 3, 2006)* (Ye. Likhoshway, ed.), pp. 115–123, Biopress, Bristol.
- Temniskova, D.N. & D.G. Vodenicharov 1981. Electron microscopic investigations on *A. praeislandica* (Jousé) Sim. descr. emend. Temn. et Voden. *Comptes rendus de L'Académie bulgare des Sciences* 34(8):1127–1130.
- Tsoy, I.B. 2017. Early Miocene diatom flora from the Yamato Rise, the Sea of Japan. *Diatom Research* 32(3):277–293.
- Tsoy, I.B. & M.V. Usoltseva 2015. Early Miocene freshwater diatoms from the Krishtofovich Rise (Ulleung Plateau) sediments of the Japan Sea. *Voprosy Sovremennoi Algologii* 3(10):66 (in Russian with English summary). [Цой И.Б., Усольцева М.В. 2015. Раннемиоценовые пресноводные диатомеи из отложений возвышенности Криштофовича (плато Уллын) Японского моря // Вопросы современной альгологии. № 3 (10). С. 66].
- Tsuchi, R. & J.C. Ingle 1992. *Pacific Neogene environment, evolution and events*. University of Tokyo Press, Tokyo, 257 pp.
- Usoltseva, M.V., A. Hassan, E.V. Rodionova, L.A. Titova, I.S. Chuvashova & S.V. Rasskazov 2020. The first finding of diatoms from the Early Miocene lacustrine deposits of the Barguzin Valley (Baikal Rift Zone). *Limnology and Freshwater Biology* (4):752–754.
- Usoltseva, M.V. & G.K. Khursevich 2013. *Alveolophora robusta* comb. nov. from Miocene deposits of the Vitim Plateau, Russia. *Diatom Research* 28(1):109–114.
- Usoltseva, M., G. Khursevich, S. Rasskazov, S. Vorob'eva & G. Chernyaeva 2010. Morphology of *Actinocyclus* and *Lobodiscus* species (Bacillariophyta) from the Miocene deposits of the Vitim Plateau, Russia. *Plant Ecology and Evolution* 143(3):1–13.
- Usoltseva, M., O. Likhacheva, V. Pushkar & A. Avramenko 2018. *Alveolophora khursevichiae* sp. nov. from the Miocene sediments of Khanka Lake (Far East). *Nova Hedwigia* 147:27–33.
- Usoltseva, M., O. Likhacheva, Yu. Dubrovina, V. Pushkar & E. Elbakidze 2011. Valve ultrastructure of dominant species (Bacillariophyta) from Middle Miocene Novokachalinskaya suite (Primorye). In: *Abstract of the 5th Central European Diatom Meeting (Szczecin, Poland, March 24–27, 2011)* (M. Bak, ed), p. 57, Wydawnictwo Naukowe Uniwersytetu Szczecińskiego, Szczecin, Poland.
- Usoltseva, M.V. & E.V. Likhoshway 2005. Restudy of the type material of diatom genus *Aulacoseira* from the Miocene deposits of Pra-Khanka Lake by scanning electron microscopy. In: *Proceeding of the First all-russian school "Modern Paleontology: classic and newest methods"* (A.Yu. Rozanov, A.V. Lopatin, P.Yu. Parkhaev, eds.), pp. 181–186, ARES, Moscow (in Russian with English summary). [Усольцева М.В., Лихошвай Е.В. 2005. Изучение типового материала диатомовых водорослей рода *Aulacoseira* из миоценовых отложений озера пра-Ханка с помощью сканирующей электронной микроскопии // Материалы I Всероссийской научной школы молодых ученых-палеонтологов «Современная палеонтология: классические и новейшие методы» (Москва, 20–24 октября 2004 г.) / отв. ред. А.Ю. Розанов, А.В. Лопатин, П.Ю. Пархаев. Москва: АРЕС. С. 181–186].
- Usoltseva, M.V. & E.V. Likhoshway 2007. The fine structure of loricae in the species of the genus *Aulacoseira* Thwaites (Bacillariophyta) from Ob' River (Russia). *Algologiya* 17(2): 139–147 (in Russian with English summary). [Усольцева М.В., Лихошвай Е.В. 2007. Тонкое строение панцирей видов рода *Aulacoseira* Thwaites (Bacillariophyta) из реки Обь // Альгология. Т. 17, № 2. С. 139–147].
- Usoltseva, M.V. & I.B. Tsoy 2010. Elliptical species of the freshwater genus *Aulacoseira* in Miocene sediments from Yamato Rise (Sea of Japan). *Diatom Research* 25(2):397–415.
- Zachos, J., M. Pagani, L. Sloan, E. Thomas & K. Billups 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693.
- Wang, P., Q. Zhao, Z. Jian, X. Cheng, W. Huang, J. Nian, J. Wang, Q. Li, B. Li & X. Sun 2003. Thirty million year deep-sea records in the South China Sea. *Chinese Science Bulletin* 48(23):2524–2535.
- Zhisheng, A. 2000. The history and variability of the East Asian paleomonsoon climate. *Quaternary Science Reviews* 19:171–187.
- Zhisheng, A., J.E. Kutzbach, W.L. Prell & S.C. Porter 2001. Evolution of Asian monsoons and phased uplift of the Himalaya-Tibetan plateau since Late Miocene times. *Nature* 411:62–66.